



World Health Organization

Western Pacific Region



Volume 4, Number 4, 2013, Pages 1–29
p-ISSN: 2094-7321 e-ISSN: 2094-7313

Perspective

Efficient use of social media during avian influenza A(H7N9) emergency response 1

Fung I and Wong K

Surveillance Report

Invasive meningococcal disease in elderly people, New South Wales, Australia, 1993 to 2012 4

Gunaratnam P, Massey P, Durrheim D, Torvaldsen S

Surveillance of avian influenza viruses in Papua New Guinea poultry, June 2011 to April 2012 11

Jonduo MH, Wong SS, Kapo N, Omonipi P, Abdad MY, Siba PM, McKenzie P, Webby R, Horwood P

Original Research

Surveillance for viral zoonoses in New Zealand birds 16

Tompkins DM, Jakob-Hoff R, Pullford D, Castro I, Mackereth G

Letter to the Editor

Letter to the Editor: Human infections with avian influenza A(H7N9) virus in China: preliminary assessments of the age and sex distribution 24

Wiwanitkit V

Author Response

Author response: Human infections with avian influenza A(H7N9) virus in China: preliminary assessments of the age and sex distribution 24

Arima Y, Zu R, Murhekar M, Vong S and Shimada T

Western Pacific Surveillance and Response Journal instructions to authors 26

Acknowledgements to WPSAR reviewers 2013 29

Western Pacific Surveillance and Response

Open access journal with continuous publication

Western Pacific Surveillance and Response (WPSAR) is an open access journal dedicated to the surveillance of and response to public health events. The goal of the journal is to create a platform for timely information sharing both within our region and globally to enhance surveillance and response activities. WPSAR is a continuous publication which means articles will be published online as soon as they have completed the review and editing process. Every three months articles will be batched for a print issue. It is a publication managed by the World Health Organization Regional Office for the Western Pacific.



EDITORIAL TEAM

Ailan Li
Executive Editor

Michelle McPherson
Coordinating Editor

Elizabeth Mangali
Assistant Editor

Associate Editors

Jorge Mendoza Aldana

Frank Konings

Nobuyuki Nishikiori

Manju Rani

Boris Pavlin

Dongbao Yu

Copyright notice

© World Health Organization 2010

p-ISSN: 2094-7321

e-ISSN: 2094-7313

All rights reserved. The information presented in the various pages of this journal is issued by the World Health Organization for general distribution, and is protected under the Berne Convention for the Protection of Literature and Artistic Works, under national laws on copyright and neighbouring rights.

The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use.

Publications of the World Health Organization can be obtained from Marketing and Dissemination, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; email: bookorders@who.int). Requests for permission to reproduce WHO publications, in part or in whole, or to translate them – whether for sale or for non-commercial distribution – should be addressed to Publications, at the above address (+41 22 791 4806; e-mail: permissions@who.int). For WHO Western Pacific Regional Publications, request for permission to reproduce should be addressed to Publications Office, World Health Organization, Regional Office for the Western Pacific, P.O. Box 2932, 1000, Manila, Philippines, fax: +632 521 1036, e-mail: publications@wpro.who.int.

Disclaimer

The designations employed and the presentation of the information in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

To contact us:

Western Pacific Surveillance and Response

World Health Organization
Office for the Western Pacific Region
United Nations Avenue
1000 Manila, Philippines
wpsar@wpro.who.int
www.wpro.who.int/wpsar

Efficient use of social media during the avian influenza A(H7N9) emergency response

Isaac Chun-Hai Fung,^{ab} and Karen K Wong,^c

Correspondence to Isaac Chun-Hai Fung (e-mail: CFung@georgiasouthern.edu).

During the 2013 outbreak of human infections of avian influenza A(H7N9), the Centers for Disease Control and Prevention (CDC) used official data released by the World Health Organization (WHO) and the Chinese government to keep United States public health officials informed of updates of the outbreak.¹ The Chinese central government released official avian influenza A(H7N9) data via its web sites (e.g. National Health and Family Planning Commission²), their official news agency (Xinhua News Agency) and their official newspapers (e.g. *People's Daily*, Beijing). In addition, official avian influenza A(H7N9) information was released by Chinese provincial and municipal governments such as Shanghai Municipal Bureau of Health,³ Jiangsu Department of Health⁴ and Zhejiang Department of Health.⁵ Prior studies have discussed the role of social media in the early detection of disease outbreaks^{6–9} and the facilitation of community-level discussion.¹⁰ In this perspective, we focus on the use of social media by public health agencies to disseminate and obtain official outbreak information during a public health emergency response.

Weibo (literally, microblog) is a category of Chinese microblogging sites that are similar to Twitter. Both Twitter and *weibo* are social media that allow users to post a 140-character long message online. *Weibo* has become popular in China since August 2009 when Twitter became unavailable to users in mainland China. As of December 2012, 309 million people were reported to be *weibo* users in China as compared to the global 500 million registered Twitter users as of July 2012. There are several different providers of *weibo*, including *Sina Weibo*, *Tencent (QQ) Weibo*, *Sohu*

Weibo, *Baidu Weibo*, *ifeng Weibo*, *NetEase Weibo* and others. Most *weibo* users live in China; a random sample of users of *Sina Weibo* found that 1.6% of users were from countries other than China.¹¹

Social media platforms provide a new channel through which public health agencies release official information, either by posting new outbreak information directly or by guiding people to official web sites. The 2013 H7N9 outbreak was the first time that WHO used Twitter for initial release of official outbreak information.¹² Likewise, the Chinese central government, some of its provincial and municipal governments and the Chinese official news agency released some official outbreak information via *weibo* nearly simultaneously with their web site press releases (the exact time of information release is known for *weibo* but often not for web sites; [Table 1](#)). An official list of Chinese provincial and municipal health authorities' *weibo* accounts can be found at the web site of the National Health and Family Planning Commission.¹³ Social media, like Twitter and *weibo*, are used by WHO and the Chinese authorities to direct attention of online communities towards their official web site press releases ([Table 1](#)). *Weibo* users can also post text longer than 140 characters as an image attached to their *weibo* post, which is known as a long *weibo*. The Chinese government used this function to post press releases on *weibo*. An example of a long *weibo* post containing a whole press release by the Shanghai Municipal Government¹⁴ can be found in [Table 1](#).

Social media platforms can help CDC epidemiologists obtain official information more efficiently because

^a Health Economics and Modeling Unit, Division of Preparedness and Emerging Infections, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, United States of America.

^b Department of Epidemiology, Jiann-Ping Hsu College of Public Health, Georgia Southern University, Statesboro, Georgia, United States of America.

^c Epidemic Intelligence Service assigned to Influenza Division, National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, United States of America.

Submitted: 17 September 2013; Published: 31 October 2013

doi: 10.5365/wpsar.2013.4.3.005

Table 1. Examples of outbreak information released online and through social media by the World Health Organization and the Chinese national, provincial and municipal health authorities

Organization	Social media/web site	Title and content (web site address)	Date/time (if available)	Language
World Health Organization (WHO)	@WHO	China has notified WHO of three human cases infected with #influenza A(#H7N9) http://goo.gl/Wf7Fy #Flufighter (https://twitter.com/WHO/status/318764531029516288)	1 April 2013 16:39:21 GMT	English
	http://www.who.int/	Human infection with influenza A(H7N9) virus in China. Global Alert and Response – Press release (http://www.who.int/csr/don/2013_04_01/en/index.html)	1 April 2013*	English
Chinese central government – National Health and Family Planning Commission	http://e.weibo.com/u/2834480301 (健康中国)	卫生和计划生育委员会就上海、安徽3例患者感染H7N9禽流感展开疫情问答 http://t.cn/zTwy8Jw 确诊患者的所有密切接触者目前未发现类似病例。 [Translated: The National Health and Family Planning Commission provides a Q&A session on epidemiologic information regarding the three patients infected with H7N9 avian influenza in Shanghai and Anhui (http://t.cn/zTwy8Jw) up to the present; there are no similar cases found among all the close contacts of the confirmed cases (http://e.weibo.com/2834480301/zq0QvBVsl).]	31 March 2013 11:38 GMT	Chinese
	http://www.moh.gov.cn/	上海、安徽发生3例人感染H7N9禽流感确诊病例 [Translated: three confirmed cases of human infections of H7N9 avian influenza in Shanghai and Anhui – Press release] (http://www.moh.gov.cn/mohwsyjbg/s3578/201303/44f25bd6bed14cf082512d8b6258fb3d.shtml)	31 March 2013*	Chinese
Chinese provincial/municipal government – e.g. Shanghai Municipal Government	http://e.weibo.com/shanghai (上海发布)	【上海两例人感染H7N9禽流感病例密切接触者未发现异常情况】#要闻#经国家卫生和计划生育委员会组织专家确诊并依法向社会公布，上海市发现2例人感染H7N9禽流感病例。截至目前，所有密切接触者均未发现类似症状和发病情况。今年以来上海流感、肺炎发病率与近三年同期相比并无明显上升。详见长微博。 [Translated: (Two cases of human infection of H7N9 avian influenza in Shanghai. Abnormal situation has not been observed among close contacts.) #Headline# The National Health and Family Planning Commission has organized experts to confirm and has followed the law to report to the society, that two cases of human infection of avian influenza have been discovered in Shanghai. Up to the present, there has not been any similar symptoms or disease onset observed among all close contacts. Since the beginning of this year, the clinical attack rates of influenza and pneumonia in Shanghai are similar to the same period in the past three years and no obvious increase has been observed. For details, see long <i>weibo</i> post.] (http://e.weibo.com/2539961154/zq0AU9VMR)	31 March 2013 11:00 GMT	Chinese
	http://wsj.sh.gov.cn/ http://www.smhb.gov.cn/	上海、安徽发生3例人感染H7N9禽流感确诊病例 [Translated: three confirmed cases of human infections of H7N9 avian influenza in Shanghai and Anhui – Press release] (http://wsj.sh.gov.cn/website/b/103667.shtml)	31 March 2013*	Chinese

* The precise release time for the official press releases was not available as the webpages did not carry a stamp of their release time. Nonetheless, based on our experience, the online press releases and the social media posts were released by WHO and the Chinese authorities nearly simultaneously.

information from multiple sources can be obtained from a central access point. During the avian influenza A(H7N9) outbreak, a team at CDC followed the social media accounts of multiple official sources so that new outbreak information from WHO and the Chinese health authorities would automatically come to the team's attention. When new case data were released by WHO or the Chinese government at its national,

provincial or municipal level via Twitter and/or *weibo*, the event was re-tweeted by social media users; thus even a message originating on a Chinese-language web site of a provincial health department would be rapidly noticed worldwide and quickly rise to the team's attention. While these social media posts might include information that was already available elsewhere, they did alert epidemiologists to the release of new

information through official sources, allowing the team to gather additional information from official web sites if available (Table 1) and obviating the need for constant monitoring of multiple news sources and web sites, such as individual web sites of the many local Chinese health departments.

The use of Chinese social media, like *weibo*, coupled with the necessary Chinese language and cultural knowledge, enabled CDC epidemiologists to gather the Chinese official data so that it could be translated, contextualized and interpreted in an efficient manner during the A(H7N9) emergency response. To ensure timely and complete understanding of an outbreak situation, it may be helpful for epidemiologists to track social media, including Twitter and *weibo*, in addition to traditional methods of communication.⁷ Our experiences in the 2013 avian influenza A(H7N9) outbreak could be relevant to other outbreaks in other countries and to public health agencies of other nations.

Disclaimer

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the CDC.

Conflicts of interest

None declared.

Funding

None

Acknowledgement

We want to thank the team members: Fiona Havers, Su Su, Kira Christian and the Global Disease Detection Operations Center, Center for Global Health. We thank Fiona Havers, Wenkai Li, Martin I Meltzer, Scott Santibanez and Michael Washington for their comments on earlier versions of this manuscript.

References

- Centers for Disease Control and Prevention. Emergence of Avian Influenza A(H7N9) Virus Causing Severe Human Illness — China, February–April 2013. *Morbidity and Mortality Weekly Report*, 2013, 62(18):366–371. pmid:23657113
- National Health and Family Planning Commission. Beijing, 2013 (<http://www.moh.gov.cn/>, accessed 21 August 2013).
- Shanghai Municipal Bureau of Health. Shanghai, 2013 (<http://wsj.sh.gov.cn/>; <http://www.smhb.gov.cn/>, accessed 21 August 2013).
- Jiangsu Department of Health. Jiangsu, 2013 (<http://www.jsbst.gov.cn/>, accessed 21 August 2013).
- Zhejiang Department of Health. Zhejiang, 2013 (<http://www.zjwst.gov.cn/>, accessed 21 August 2013).
- Salathé M, Khandelwal S. Assessing vaccination sentiments with online social media: implications for infectious disease dynamics and control. [Validation Studies]. *PLoS Computational Biology*, 2011, 7:e1002199. doi:10.1371/journal.pcbi.1002199 pmid:22022249
- Salathé M et al. Influenza A (H7N9) and the importance of digital epidemiology. *The New England Journal of Medicine*, 2013, 369:401–404. doi:10.1056/NEJMp1307752 pmid:23822655
- Chew C, Eysenbach G. Pandemics in the age of Twitter: content analysis of Tweets during the 2009 H1N1 outbreak. *PLoS ONE*, 2010, 5:e14118. doi:10.1371/journal.pone.0014118 pmid:21124761
- Signorini A, Segre AM, Polgreen PM. The use of Twitter to track levels of disease activity and public concern in the U.S. during the influenza A H1N1 pandemic. *PLoS ONE*, 2011, 6:e19467. doi:10.1371/journal.pone.0019467 pmid:21573238
- Chan M et al. Microblogging, online expression, and political efficacy among young Chinese citizens: the moderating role of information and entertainment needs in the use of *Weibo*. *Cyberpsychology, behaviour and social networking*, 2012, 15(7):345–349.
- Fu KW, Chau M. Reality check for the Chinese microblog space: a random sampling approach. *PLoS*, 2013, 8:e58356. doi:10.1371/journal.pone.0058356 pmid:23520502
- From now on Twitter will be our first place for posting #H7N9 case updates. Full updates to follow on our website as usual. Stay tuned! Geneva, World Health Organization, 2013 (<https://twitter.com/WHO/status/320148152499982336>, accessed 21 August 2013).
- Weibo matrix*. Beijing, National Health and Family Planning Commission, 2013 (<http://www.moh.gov.cn/zhuzhan/wbjz/weibo.shtml>, accessed 16 October 2013).
- Two cases of human infection of H7N9 avian influenza in Shanghai. Abnormal situation has not been observed among close contacts. Shanghai, Shanghai Municipal Government, 2013 (<http://e.weibo.com/2539961154/zq0AU9VMR>, accessed 16 October 2013).

Invasive meningococcal disease in elderly people, New South Wales, Australia, 1993 to 2012

Praveena Gunaratnam,^{ab} Peter Massey,^{c,e} David Durrheim^{cf} and Siranda Torvaldsen^b

Correspondance to Praveena Gunaratnam (e-mail: pguna@health.nsw.gov.au).

Little information is available publicly on invasive meningococcal disease (IMD) in elderly people in Australia. This study analysed IMD notifications data from New South Wales between 1993 and 2012 to determine the distribution of IMD among people aged 65 years and older and to describe the characteristics of IMD in this age group compared to younger age groups with respect to notification trends, serogroup distribution and mortality rates. Following introduction of a childhood vaccination programme against meningococcal type C in 2003, notification rates in all age groups decreased, but the proportion of IMD notifications in people aged 65 years and over rose significantly (from 4% to 6%, $P = 0.01$). Mortality rates from IMD in those aged 65 years and older were significantly higher than overall rates (32% compared to 5%, $P < 0.01$). Serogroup Y accounted for 23% of infections in the elderly compared to 3% in people aged under 65 years ($P < 0.01$). As the population ages, the elderly may account for a higher number of IMD cases in Australia. Protocols at the state and national level should be updated to provide guidance on the clinical and public health management of elderly people with IMD.

Invasive disease caused by *Neisseria meningitidis* occurs when bacteria enter a normally sterile site such as blood (causing septicaemia) or cerebrospinal fluid (causing meningitis).¹ Transmission is by respiratory droplets and up to 10% of the population may harbour *Neisseria meningitidis* in their nasopharynx without disease.¹ Asymptomatic carriage is higher in household contacts of patients with meningococcal disease (12.4%).² In Australia, the highest rates of invasive meningococcal disease (IMD) are among children under 5 years and young adults between 15 and 24 years of age.³

Five serogroups (A, B, C, W135 and Y) are responsible for the majority of IMD worldwide,⁴ with serogroups B and C the most commonly reported in Australia.¹ Serogroup C vaccine is the only meningococcal vaccine routinely offered to all children in Australia under the National Immunization Programme. The national meningococcal C vaccination programme, introduced in Australia in January 2003, comprises ongoing inclusion of the vaccine in the National Immunization Programme schedule at one year of age, as well as a catch-up programme until 2006 for children aged 2–19 years.^{3,4} Large decreases have since been observed in serogroup C infections,

with a 92% decline in the number of notifications of this serotype in the 15–24 year age group.⁴ Across all age groups, notification of IMD in Australia decreased by half from a rate of 2 per 100 000 to 1 per 100 000 population between 2004 and 2010.¹

The state of New South Wales (NSW; the largest Australian state) has a population of over 7.2 million; 1 million (14%) are aged 65 years and over. The population is ageing, and the number of people in this age group grew by 22% between 2001 and 2011.⁵ Throughout Australia, the proportion of the population aged 65 years and over is projected to increase to just under a quarter of the total population by 2056.⁶ Confirmed and probable cases of IMD in NSW have been notifiable by clinicians and laboratories since 1990 under the NSW Public Health Act (**Box 1**). Case notification and surveillance data are entered into the NSW Notifiable Conditions and Information Management System, including details on age and serogroup information where available.⁷

Little has been reported on IMD incidence in the elderly internationally, with no reports published in Australia. In late 2012, the Hunter New England Local

^a New South Wales Public Health Officer Training Program, New South Wales Health, Sydney, Australia.

^b School of Public Health and Community Medicine, University of New South Wales, Sydney, Australia.

^c Hunter New England Population Health, Newcastle, Australia.

^d School of Health, University New England, Armidale, Australia.

^e School of Nursing, Midwifery & Nutrition, James Cook University, Townsville, Australia.

^f Hunter Medical Research Institute, New Lambton Heights, Australia.

Submitted: 30 September 2013; Published: 9 December 2013

doi: 10.5365/wpsar.2013.4.4.001

Box 1. Meningococcal disease case definitions, New South Wales, Australia

Case	Definition
Confirmed	Requires either laboratory definitive evidence, OR laboratory suggestive evidence AND clinical evidence. Laboratory definitive evidence - Isolation of <i>Neisseria meningitidis</i> from a normally sterile site; OR - Detection of specific meningococcal DNA sequences in a specimen from a normally sterile site by nucleic acid amplification testing. Laboratory suggestive evidence - Detection of Gram negative diplococci in Gram stain of specimen from a normally sterile site or from a suspicious skin lesion; OR - High titre IgM or significant rise in IgM or IgG titres to outer membrane protein antigens of <i>Neisseria meningitidis</i>. Clinical evidence - A disease which is compatible with invasive meningococcal disease in the opinion of the treating clinician.
Probable	Requires clinical evidence only: - The absence of evidence for other causes of clinical symptoms; AND EITHER - Clinically compatible disease including haemorrhagic rash; OR - Clinically compatible disease AND close contact with a confirmed case within the previous 60 days.

Health District in northern NSW was notified of a case of IMD in a resident of an aged care facility. The response to this case revealed a lack of published guidance with respect to the public health management of IMD in the elderly, particularly in the definition of close contacts and administration of clearance treatment.

The aim of the current epidemiological investigation was to examine IMD trends in NSW with a particular focus on the proportion of notifications in people aged 65 years and over, mortality rates and serogroup distribution. Implications for the prevention and management of IMD are also discussed.

METHODS

IMD notification data for the period 1993 to 2012 for NSW were sourced from the NSW Ministry of Health. Analysis was performed using Microsoft Excel, Statistical Analysis Software (SAS) Enterprise Guide 5.1 (SAS Institute Inc. Seattle, WA, USA) and STATA 11 (Stata Corp. 2009, College Station, TX, USA). IMD notification rates were calculated using mid-year NSW population data from the Australian Bureau of Statistics and proportions were compared using the chi-squared (χ^2) test.

Annualized notification rates by age groups (10–14, 0–14, 15–24, 25–64 year olds and people aged 65 years and over) were compared over two periods, 1993 to 2002 and 2003 to 2012, corresponding to periods before and after introduction of the national childhood meningococcal C vaccination programme in 2003. The 0–14 year age group was further divided into 0–4, 5–9 and 10–14 years. Wider age ranges were

chosen to facilitate comparison between people aged 65 years and over and younger age groups, particularly those aged under 25 who are considered most at risk of IMD. Rate ratios and 95% confidence intervals were calculated for each age group to compare 2003 to 2012 notification rates with 1993 to 2002.

Analysis by serogroup was conducted for 2003 to 2012, the years for which serogroup information was recorded for more than 70% of notifications; cases diagnosed by polymerase chain reaction were not further typed. Case fatality rates (CFRs) were also limited to 2008 and 2012 due to missing data before this time.

This project used de-identified data routinely collected under the NSW Public Health Act. The project was further deemed a quality improvement exercise and an ethics waiver was granted by the Hunter New England Local Health District Human Research Ethics Committee.

RESULTS

Notifications by age group

Between 1993 and 2012, there were 2995 notifications of IMD in NSW. Annual notifications peaked at 253 in 2000, decreasing to 66 in 2012 (**Figure 1**). The age group with the highest number of notifications each year was 0–14 year olds; the lowest was observed for the 65 years and over age group (**Figure 1**).

Over the 20-year period, there were 153 notifications (5.1%) in those aged 65 years and over. The

Figure 1. Invasive meningococcal disease notifications by year and age group, New South Wales, Australia, 1993 to 2012




Table 1. Changes in invasive meningococcal disease notification rates by age group, New South Wales, Australia, 1993 to 2002 and 2003 to 2012

Age group (years)	n	1993 to 2002		n	2003 to 2012		Rate ratio (95% confidence interval)	P-value
		%	Rate per 100 000		%	Rate per 100 000		
0–14	978	51.5	74.1	497	45.2	37.04	0.50 (0.45–0.56)	< 0.001
0–4	701	36.9	159.8	371	33.8	82.79	0.52 (0.46–0.59)	< 0.001
5–9	151	8.0	34.1	76	6.9	17.18	0.50 (0.38–0.66)	< 0.001
10–14	126	6.6	28.7	50	4.5	11.08	0.39 (0.28–0.53)	< 0.001
15–24	479	25.3	54.3	267	24.3	28.34	0.52 (0.45–0.61)	< 0.001
25–64	357	18.8	10.8	264	24.0	7.12	0.66 (0.56–0.77)	< 0.001
65 and over	82	4.3	10.2	71	6.5	7.37	0.72 (0.53–0.99)	0.045
Total	1896	100.0	30.0	1099	100.0	15.81	0.53 (0.49–0.57)	< 0.001

proportion of IMD notifications for those aged 65 years and over significantly increased from 4% in the period 1993–2002 to 6% in the period 2003–2012 ($P = 0.01$). Although annualized rates per 100 000 population decreased significantly across all age groups between 1993–2002 and 2003–2012, the decrease was primarily for younger age groups compared with those aged 65 years and over (Table 1).

Case fatality rates


Mortality data were available for 366 of 390 IMD notifications between 2008 and 2012, with 19 deaths recorded: three in 0–14 year olds, one in 15–24 year olds,

10 in 25–64 year olds and six in those aged 65 years and over. The CFR for those aged 65 years and over was significantly higher than the overall CFR (32% [6/19] compared with 5% [19/366]; $P < 0.01$) and also than the CFR for the 0–4 year old age group (1% [3/127]; $P < 0.01$), those historically at greatest risk of mortality.

Notifications by serogroup

Between 2003 and 2012, serogroup information was available for 64/71 (90%) notifications in those aged 65 years and over and for 787/1028 (77%) notifications in people under 65 years of age. During this period, there

Figure 2. Invasive meningococcal disease notifications by year and serogroup, New South Wales, Australia, 2003 to 2012



was a gradual reduction in the proportion of infections caused by serogroup C and an increase primarily in the proportion caused by serogroup B (Figure 2).

Of the 64 notifications in those aged 65 years and older, 30 (47%) were serogroup B, 10 (16%) were serogroup C, 15 (23%) were serogroup Y and 9 were (14%) serogroup W. For those under 65 years of age, 605 (77%) were serogroup B, 125 (16%) were serogroup C, 21 (3%) were serogroup Y and 35 (4%) were serogroup W. The difference in proportions attributable to serogroup Y infection between the two age groups was statistically significant ($P < 0.01$).

Across all age groups, the proportion of IMD notifications attributable to serogroup Y significantly increased from 3% (17/539) in the period 2003–2007 to 6% (19/312) in the period 2008–2012 ($P > 0.05$). For those aged under 65 years, the proportion of notifications attributable to serogroup Y experienced a slight but steady increase between 2003 and 2012, while for older age groups there was variability from year to year (Figure 3).

DISCUSSION


The absolute number of IMD cases in NSW residents aged 65 years and over has remained low over the past 20 years, and rates for all age groups have decreased,

most likely as a result of the herd immunity effect of the childhood meningococcal C vaccination programme. However, the 65 years and over age group constituted an increasing proportion of all notifications had a higher proportion of infections of serogroup Y. Almost one third of cases resulting in death.

An increase in the proportion of IMD due to serogroup Y has occurred in NSW during the past 10 years. This has been previously reported throughout Australia, although statistical significance was not demonstrated.⁸ Research in the United States of America and United Kingdom has also found a higher prevalence of serogroup Y in older people.^{9,10} Higher CFRs have previously been reported in older people with IMD (particularly among those with underlying medical conditions) and among serogroup Y cases even after controlling for age.^{9–11} This may explain the higher CFR in older persons in this study; they had a higher proportion of infection with serogroup Y.

Potential increases in the number of IMD cases among the elderly, as a result of an ageing population, have implications not only for the clinical management of these cases but also for the public health response. The current *Guidelines for the early clinical and public health management of meningococcal disease in Australia* recommend provision of clearance antibiotics to household contacts or household-like contacts in

Figure 3. Proportion of invasive meningococcal disease notifications attributable to serogroup Y by year and age group, New South Wales, Australia, 2003 to 2012



settings such as schools or universities but do not specifically address aged care or residential facilities.¹² A similar lack of specificity is also found in other national and regional guidelines (Table 2) with no particular guidelines on meningococcal disease available for the Western Pacific Region.

Outbreaks in residential aged care facilities have been reported, but specific studies are limited and provide contradictory advice.¹³ Prophylaxis was administered following a case of meningococcal disease in an aged care facility in the United States of America in 1997; a further case was then reported in a patient who had refused prophylaxis.¹⁴ A carriage study in the United Kingdom conducted after a case of meningococcal disease was diagnosed in a nursing home conversely found that no residents or caregivers had the same sero-subtype of *Neisseria meningitidis* as the index case, and the study concluded that prophylaxis was not necessary in such settings.¹⁵

The analyses on serogroup and mortality presented in this study should be interpreted with caution given the amount of missing information, particularly between 1993 and 2002 and because the estimates among older people are based on relatively small numbers. It is also likely that mortality was underestimated across all age groups, given the difficulty in determining a single cause of death. However, as IMD is a notifiable and serious disease, affected patients are likely to present in a health

-care setting and be tested and notified by a treating physician, thus the notification data should approximate incidence.

CONCLUSION

The epidemiology of IMD in NSW, Australia has changed following the introduction of the childhood vaccination programmes, with a higher proportion of infections in those aged 65 years and over attributable to serogroup Y. Similar analysis should be conducted at the national and regional levels to determine whether similar trends are occurring across Australia and other countries that have introduced childhood meningococcal C vaccination programmes. As the Australian population is ageing, there may be increases in the number of notifications of IMD among people aged 65 years and over, including cases and potentially outbreaks in aged care facilities. Such an increase would have serious implications given the higher mortality rates in this group. National protocols should be updated to provide clinical and public health guidance for this age group.

Conflict of Interest

None declared.

Funding

None.

Table 2. Recommendations on use of chemoprophylaxis in general and in aged care facilities following a case of invasive meningococcal disease – various guidelines

Country or Region	Guideline	Recommendations on use of chemoprophylaxis in general and in aged care facilities
Australia	<i>Guidelines for the early clinical and public health management of meningococcal disease in Australia.</i> Canberra, Australian Government of Health and Ageing and Communicable Diseases Network Australia, October 2007. ¹²	"The household contacts of a case, including recent visitors who have stayed overnight in the 7 days preceding the onset of the case's illness should receive clearance antibiotics and vaccination. ... Those who share the same dormitory, military barrack or hostel bunkroom as a case are, in effect, household contacts." p39 No specific mention of aged care facilities.
United Kingdom	<i>Guidance for public health management of meningococcal disease in the United Kingdom.</i> London, Health Protection Agency United Kingdom, March 2012 (http://www.hpa.org.uk/webc/hpawebfile/hpaweb_c/1194947389261).	"Chemoprophylaxis should be offered to close contacts of cases... in the following categories: (a) Those who have had prolonged close contact with the case in a household type setting during the seven days before onset of illness. Examples of such contacts would be those living and/or sleeping in the same household (including extended household)... .Prophylaxis not indicated (unless already identified as close contacts) for ... residents of nursing/residential homes." p22
United States of America	<i>Prevention and Control of Meningococcal Disease: Recommendations of the Advisory Committee on Immunization Practices.</i> Atlanta, United States Centers for Disease Control and Prevention, March 2013 (http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6202a1.htm).	"Antimicrobial chemoprophylaxis of close contacts of a patient with invasive meningococcal disease is important to prevent secondary cases. Close contacts include: (1) household members (2) child-care center contacts, and (3) anyone directly exposed to the patient's oral secretions... ... in the 7 days before symptom onset." p23 No specific mention of aged care facilities.
Canada	<i>Guidelines for the Prevention and Control of Meningococcal Disease.</i> Ottawa, Public Health Agency of Canada, May 2005 (http://www.phac-aspc.gc.ca/publicat/ccdr-rmtc/05vol31/31s1/).	"Chemoprophylaxis should be provided to close contacts" p7. Close contacts defined as including: – household contacts of a case – persons who share sleeping arrangements with a case. p3 No specific mention of nursing homes other than as potential location for organization based outbreaks. p4
New Zealand	<i>Communicable Disease Control Manual – Neisseria meningitidis invasive disease.</i> Wellington, New Zealand Ministry of Health, May 2012 (http://www.health.govt.nz/publication/communicable-disease-control-manual-2012).	Contact defined as, "Anyone who has had unprotected contact with upper respiratory tract or respiratory droplets from the case during the 7 days before onset of illness to 24 hours after onset of effective treatment." p3 "Public health follow-up is important for household contacts and contacts that have had similarly close exposure. Examples of such contacts are: • those sleeping at least one night in the same household, dormitory, military barrack, student hostel bunkroom (not residents of nursing or residential homes who sleep in separate rooms) as the case" p4
Europe	<i>Public health management of sporadic cases of invasive meningococcal disease and their contacts.</i> Stockholm, European Centre for Disease Prevention and Control, October 2010 (http://www.ecdc.europa.eu/en/publications/publications/1010_gui_meningococcal_guidance.pdf).	"Chemoprophylaxis with an antibiotic regime that eradicates carriage is recommended for household contacts of a case of IMD." p12 No definition of "household contacts" or mention of aged care facilities specifically.

Acknowledgements

Praveena Gunaratnam is employed as a trainee on the NSW Public Health Officer Training Programme, funded by the NSW Ministry of Health. She undertook this work based at the Hunter New England Population Health. The Programme is offered in partnership with the University of NSW.

References:

1. NNDSS Annual Report Writing Group. Australia's notifiable disease status, 2010. *Communicable Diseases Intelligence*, 2012, 36: 1–69.
2. Kristiansen BE, Tveten Y, Jenkins A. Which contacts of patients with meningococcal disease carry the pathogenic strain of *Neisseria meningitidis*? A population based study. *BMJ (Clinical research ed.)*, 1998, 317:621–625. doi:10.1136/bmj.317.7159.621 pmid:9727987

3. Australian Government Department of Health and Ageing and the National Health and Medical Research Council. *The Australian Immunisation Handbook 9th Edition*. Canberra, Australian Government Department of Health and Ageing, 2008 ([http://www.immunise.health.gov.au/internet/immunise/publishing.nsf/Content/78CDF41C283426A8CA2574E40020CCAB/\\$File/handbook-9.pdf](http://www.immunise.health.gov.au/internet/immunise/publishing.nsf/Content/78CDF41C283426A8CA2574E40020CCAB/$File/handbook-9.pdf), accessed 15 September 2013).
4. Halperin SA et al. The changing and dynamic epidemiology of meningococcal disease. *Vaccine*, 2012, 30(Suppl 2):B26–36. doi:10.1016/j.vaccine.2011.12.032 pmid:22178525
5. *Population by Age and Sex, Regions of Australia, 2011*. Cat No 3235.0. Canberra, Australian Bureau of Statistics, 2012.
6. *Australian Social Trends, September 2009*. Cat No 4102.0. Canberra, Australian Bureau of Statistics, 2009.
7. *Control Guideline: Meningococcal Disease, last updated: 01 July 2012*. Australia, New South Wales Health, 2012 (<http://www.health.nsw.gov.au/Infectious/controlguideline/Pages/mening.aspx>, accessed 15 September 2013).
8. Lahra MM, Enriquez RP. Annual report of the Australian Meningococcal Surveillance Programme, 2011. *Communicable Diseases Intelligence*, 2012, 36:E251–62. pmid:23186236
9. Cohn AC et al. Changes in *Neisseria meningitidis* disease epidemiology in the United States, 1998–2007: implications for prevention of meningococcal disease. *Clinical Infectious Diseases*, 2010, 50:184–191. doi:10.1086/649209 pmid:20001736
10. Ladhani SN et al. Invasive meningococcal capsular group Y disease, England and Wales, 2007–2009. *Emerging Infectious Diseases*, 2012, 18:63–70. doi:10.3201/eid1801.110901 pmid:22261040
11. Centers for Disease Control and Prevention. Prevention and Control of Meningococcal Disease: Recommendations of the Advisory Committee on Immunization Practice. *Morbidity and Mortality Weekly Report*, 2013, 62(2) (<http://www.cdc.gov/mmwr/pdf/rr/rr6202.pdf>, accessed 15 September 2013).
12. Australian Government Department of Health and Ageing and Communicable Diseases Network Australia. *Guidelines for the early clinical and public health management of meningococcal disease in Australia*. Canberra, Australian Government Department of Health and Ageing, 2009 ([http://www.health.gov.au/internet/main/publishing.nsf/Content/4DFF673115F66413CA257BF00020630F/\\$File/meningococcal-guidelines.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/4DFF673115F66413CA257BF00020630F/$File/meningococcal-guidelines.pdf), accessed 15 September 2013).
13. Brooks R et al. Increased case-fatality rate associated with outbreaks of *Neisseria meningitidis* infection, compared with sporadic meningococcal disease, in the United States, 1994–2002. *Clinical Infectious Diseases*, 2006, 43:49–54. doi:10.1086/504804 pmid:16758417
14. Centers for Disease Control and Prevention. Outbreaks of group B meningococcal disease—Florida, 1995 and 1997. *Morbidity and Mortality Weekly Report*, 1998, 47:833–837. pmid:9780239
15. Irwin DJ et al. Carriage of *Neisseria meningitidis* in residents and staff at a residential home for elderly people following a case of invasive disease. *Communicable Disease and Public Health*, 2000, 3:172–174. pmid:11014028

Surveillance of avian influenza viruses in Papua New Guinean poultry, June 2011 to April 2012

Marinjho Jonduo,^a Sook-San Wong,^b Nime Kapo,^c Paskalis Ominipi,^c Mohammad Abdad,^a Peter Siba,^a Pamela McKenzie,^b Richard Webby^b and Paul Horwood^a

Correspondence to Paul Horwood (paul.horwood@pngimr.org.pg).

We investigated the circulation of avian influenza viruses in poultry populations throughout Papua New Guinea to assess the risk to the poultry industry and human health. Oropharyngeal swabs, cloacal swabs and serum were collected from 537 poultry from 14 provinces of Papua New Guinea over an 11-month period (June 2011 through April 2012). Virological and serological investigations were undertaken to determine the prevalence of avian influenza viruses. Neither influenza A viruses nor antibodies were detected in any of the samples. This study demonstrated that avian influenza viruses were not circulating at detectable levels in poultry populations in Papua New Guinea during the sampling period. However, avian influenza remains a significant risk to Papua New Guinea due to the close proximity of countries having previously reported highly pathogenic avian influenza viruses and the low biosecurity precautions associated with the rearing of most poultry populations in the country.

Influenza virus is a major respiratory pathogen that infects an average of 5–15% of the global population each year, with approximately 500 000 human deaths related to influenza annually.¹ Currently all known influenza A viruses are naturally maintained in aquatic birds.² Occasionally these influenza viruses of avian lineage cross natural species barriers and infect other susceptible bird species and/or mammals including humans, pigs and horses. The interspecies transmission of highly pathogenic avian influenza (HPAI) virus to poultry populations often results in devastating disease outbreaks.

In 1996, a HPAI strain of H5N1 emerged in South-East Asia and extended throughout several Asian, Middle Eastern, African and European countries. Its re-emergence in 2003 resulted in the death of more than 62 million birds in Thailand alone, almost half of which were backyard poultry.³ Death caused by infection and preventive measures (such as depopulation) implemented to control the spread of the HPAI H5N1 virus resulted in considerable socioeconomic burdens for many of the affected countries.⁴ The recent emergence of a novel H7N9 virus in China (March 2013) has increased fears about the spread of influenza viruses with pandemic potential from poultry populations.⁵ The transmission

of these viruses over long distances by migrating birds is a concern for countries such as Papua New Guinea that have large poultry populations with few biosecurity precautions.

Poultry production accounts for 45% of the total annual livestock production in Papua New Guinea, and poultry consumption is second only to pigs.⁶ The short turn-around time, ease in rearing, market demand and high income from poultry production makes it more profitable than most other livestock rearing in Papua New Guinea. Most poultry farming in the country is conducted in semi-enclosed areas or free-ranged village settings. Relatively few poultry farms are commercialized and therefore do not have high biosecurity settings to reduce potential introduction of influenza viruses into the poultry population. The free-ranged village/backyard chickens are often raised together with other animals within the same pen (e.g. pigs and ducks). The village chickens also have unrestricted access to water and feed sources that may be utilized by wild birds, thus increasing the risk of exotic disease transmission.

In this paper we report a cross-sectional study to determine the presence of circulating avian influenza

^a Papua New Guinea Institute of Medical Research, Goroka, Papua New Guinea.

^b Center of Excellence for Influenza Research and Surveillance, St Jude Children's Research Hospital, Memphis, United States of America.

^c Papua New Guinea National Agriculture Quarantine and Inspection Authority, Port Moresby, Papua New Guinea.

Submitted: 21 October 2013; Published: 19 December 2013

doi: 10.5365/wpsar.2013.4.4.004

Table 1. Summary of the poultry* sampling sites in Papua New Guinea

Sampling site (Town, Province)	Number of sub-sites	Biosecurity classification			Total
		Low	Medium	High	
Daru, Western Province	18	69 (13)	0	43	112 (13)
Goroka, Eastern Highlands Province	5	25	28 (9)	0	53 (9)
Mt Hagen, Western Highlands Province	6	15 (3)	20 (2)	24	59 (5)
Mendi, Southern Highlands Province	2	0	6	0	6
Lae, Morobe Province	4	27 (4)	36 (8)	25	88 (12)
Kavieng, New Ireland Province	7	20	8	0	28
Port Moresby, Central Province	4	8	14 (5)	0	22 (5)
Madang, Madang Province	1	0	22 (9)	0	22 (9)
Rabaul, East New Britain Province	6	10	10 (2)	0	20 (2)
Kimbe, West New Britain Province	8	25	5	2	32
Vanimo, West Sepik Province	1	20 (7)	20	0	40 (7)
Kundiawa, Simbu Province	1	2	2	0	4
Wabag, Enga Province	8	6 (2)	12	0	18 (2)
Alotau, Milne Bay Province	11	15	17 (6)	0	32 (6)
TOTAL	82	242 (29)	200 (41)	94	536 (70)

* Samples in brackets were from ducks (unknown species) with the remaining from chickens.

viruses and the seroprevalence of neutralizing antibodies to avian influenza viruses in poultry populations across Papua New Guinea.

MATERIALS AND METHODS


Oropharyngeal swabs, cloacal swabs and serum were obtained from 536 poultry (466 chickens and 70 ducks) from 82 sub-sites within 14 selected provinces from June 2011 to April 2012 ([Table 1](#) and [Figure 1](#)). Qualified field officers from the Papua New Guinea National Agriculture Quarantine and Inspection Authority carried out the sampling during their routine surveillance programme, adhering to the guidelines of the Food and Agriculture Organization of the United Nations (FAO) for avian sampling.⁷

Sampling was conducted in three types of biosecurity settings: high, medium and low. These classifications were based on the amount of exposure the sampled poultry population had to other birds and/or animals. Thus, poultry sites with little-to-no exposure to other animals or birds were classified as high (e.g. commercial farms); sites with some exposure were classified as medium (e.g. semi-enclosed farms); and sites with unlimited exposure were classified as low biosecurity containment (e.g. free-range village chickens).

Oropharyngeal swabs, cloacal swabs and serum were obtained from poultry and sent at 4 °C to the laboratory for analysis. Upon arrival at the laboratory, the samples were stored at –80 °C (–20 °C for sera) until required for analysis. Total RNA was extracted from oropharyngeal and cloacal swabs using the QIAamp Viral RNA Minikit (Qiagen, Hilden, Germany), according to the manufacturer's instructions. The extracted RNA was tested for the presence of influenza A virus by real time reverse-transcriptase polymerase chain reaction (PCR) assays supplied by the Centers for Disease Control and Prevention (Atlanta, GA, USA). Samples positive or equivocal for avian influenza viruses were further tested for influenza A/H5 and A/H7 using previously published assays.⁸ Aliquots of all samples were sent to the Center of Excellence for Influenza Research and Surveillance, St Jude Children's Research Hospital (Memphis, TN, USA) for isolation and subtyping of avian influenza virus isolates.

A total of 36 paired oropharyngeal and cloacal samples collected from farms and provinces that had samples deemed equivocal were passaged three times in 10-day old embryonated chicken eggs. A sample was considered negative for isolation if no virus was isolated upon three passages. For increased sensitivity in detection of viral genome, deep-sequencing was also

Figure 1. Map of Papua New Guinea showing the 14 provinces where sampling was conducted



performed on the equivocal samples. Briefly, viral RNA was extracted, transcribed to cDNA and subjected to whole-genome amplification according to previously published methods.⁹ The resulting PCR products were then library-prepped and sequenced on the Illumina MiSeq platform (Illumina, San Diego, CA, USA) using the paired-end sequencing chemistry. After removal of MiSeq indices, analysis was performed using CLC Genomics Workbench 6.5 (CLC bio, Aarhus, Denmark) using the following process: for quality trimming sequence reads were filtered at the quality-limit threshold of 0.05; short reads and reads with more than two ambiguous bases were removed. Remaining reads were then de novo assembled using the fast-contig mapping mode at the minimum contig length of 200 base pairs; paired-reads were aligned using the scaffold option. Assembled contigs were then subjected to BLASTn search against the National Center for Biotechnology Information (Bethesda, MD, USA) database for viral sequences.

Sera were analysed for the presence of influenza A virus antibodies using the IDEXX AI MultiS Screen ELISA (IDEXX Laboratories, Rydalmere, Australia), according to the manufacturer's instructions. All serum samples were individually tested on three separate occasions to ensure the validity of results.

RESULTS

Influenza A virus was not detected in any of the oropharyngeal or cloacal swabs ($n=536$ each). Four samples had results recorded as equivocal as crossing-threshold values of 36–40 were detected. These samples were tested for influenza A/H5 and A/H7 using real-time PCR; however, all of the samples were negative. Further analysis of these samples using egg inoculation and next-generation sequencing at St Jude Children's Research Hospital (Memphis, TN, USA) resulted in no detection of influenza A virus.

Despite all serum samples being tested on three independent occasions, influenza A antibodies were not detected in any of the samples. Positive and negative control reactions supplied with the kits confirmed the validity of the results.

DISCUSSION

This paper is the first to investigate the presence and distribution of avian influenza viruses in poultry populations in Papua New Guinea. Influenza virus and antibodies were not detected in any of the samples, suggesting that there is low (or no) circulation of avian

influenza viruses in poultry in the country. Poultry and wild bird surveillance programmes in other countries, such as Australia and New Zealand, have also found low prevalence of circulating avian influenza viruses.¹⁰

The failure to detect avian influenza viruses in poultry does not necessarily mean that Papua New Guinea is at low risk for an outbreak. The recent detection of H5N1 in West Papua (Indonesia)¹¹ is a concern for Papua New Guinea as this region shares a land border with West Papua. The recent outbreak of Newcastle Disease virus in poultry populations in the north-west region of Papua New Guinea¹² highlights the potential for the incursion of exotic diseases into this region. Indeed the maiden outbreak of chikungunya was first detected in this region¹³ before subsequently spreading throughout much of the country.

Papua New Guinea is in close proximity to South-East Asian countries endemic for the H5N1 and H7N9 viruses.^{3,5} The spread of these viruses through the migration of waterfowl may be a potential source of incursion into non-endemic areas.¹⁴ Although wild bird surveillance studies have shown that there is a low prevalence of avian influenza viruses in Australia, and an absence of HPAI,¹⁰ avian influenza introduction from this direction is also a possibility given the nomadic migration of some duck species between Australia and Papua New Guinea.¹⁵ H5N1 has not been reported in the Pacific region since its re-emergence in 2003, despite being detected in the West Papua province of Indonesia. Previous studies have suggested that the Pacific region is protected from the incursion of HPAI influenza viruses by the uncommon migration of waterfowl across Wallace's Line.¹⁶ However, it is important that active surveillance continues so that outbreak mitigation steps can be rapidly implemented in the event of incursion of these viruses. In particular, future surveillance studies should focus on wild waterfowl and the potential for the introduction of avian influenza viruses through migration and nomadic movements of these birds.

In this study we report that there is no evidence of avian influenza circulation in Papua New Guinean poultry populations. However, there are some limitations to this study. A cross-sectional analysis for avian influenza viruses may not be sufficiently sensitive when a low prevalence of virus is circulating. The short lifespan of poultry bred for meat and the low number of samples

collected from each site may have contributed to the non-detection of avian influenza viruses and antibodies. Therefore, it is recommended that long-term sentinel surveillance should be established at sites where there is a risk of avian influenza introduction, such as sites close to border crossings and lakes used by waterfowl.

Although wild waterfowl migration routes are unlikely to be the source of exotic avian influenza introduction, the landborder with West Papua (Indonesia) and the poultry husbandry practices in Papua New Guinea mean that there is still a relatively high risk of introduction into the country. The introduction of HPAI viruses into Papua New Guinea could create a huge socioeconomic burden. Poultry provides the only source of protein consumption for many people in rural regions, and a large outbreak may have far-reaching health implications. Poor diagnostic capacity at a national level¹⁷ and limited outbreak response and mitigation capabilities may not be sufficient to contain an avian influenza outbreak.

Conflict of interest

None declared.

Funding

None.

Acknowledgements

This work was funded by the National Institute of Allergy and Infectious Diseases, the National Institutes of Health, under contract number HHSN266200700005C. Marinjho Jonduo was supported by a Partnership in Health Research Training Program, funded by Esso Highlands Papua New Guinea Limited.

References:

1. *Influenza (Seasonal) Fact Sheet No. 211*, 2009. Geneva, World Health Organization, 2009 (<http://www.who.int/mediacentre/factsheets/fs211/en/index.html>, accessed 15 August 2013).
2. Webster RG et al. Evolution and ecology of influenza A viruses. *Microbiological Reviews*, 1992, 56:152–179. PMID:1579108
3. Tiensin T et al. Highly pathogenic avian influenza H5N1, Thailand, 2004. *Emerging Infectious Diseases*, 2005, 11:1664–1672. doi:10.3201/eid1111.050608 PMID:16318716
4. Alexander DJ. An overview of the epidemiology of avian influenza. *Vaccine*, 2007, 25:5637–5644. doi:10.1016/j.vaccine.2006.10.051 PMID:17126960

5. Gao R et al. Human infection with a novel avian-origin influenza A(H7N9) virus. *The New England Journal of Medicine*, 2013, 368:1888–1897. doi:10.1056/NEJMoa1304459 pmid:23577628
6. Bourke RM, Harwood T (Eds). *Food and Agriculture in Papua New Guinea*. ANU E Press, The Australian National University, Canberra, 2009.
7. Whitworth D et al (Eds). *Wild Birds and Avian Influenza: an introduction to applied field research and disease sampling techniques*. *FAO Animal Production and Health Manual*, No. 5. Rome, Food and Agriculture Organization of the United Nations (FAO), 2007 (<ftp://ftp.fao.org/docrep/fao/010/a1521e/a1521e.pdf>, accessed 10 August 2013).
8. Spackman E et al. Development of a real-time reverse transcriptase PCR assay for type A influenza virus and the avian H5 and H7 hemagglutinin subtypes. *Journal of Clinical Microbiology*, 2002, 40:3256–3260. doi:10.1128/JCM.40.9.3256-3260.2002 pmid:12202562
9. Zhou B et al. Single-reaction genomic amplification accelerates sequencing and vaccine production for classical and swine origin human influenza A viruses. *Journal of Virology*, 2009, 83:10309–10313. doi:10.1128/JVI.01109-09 pmid:19605485
10. Hansbro PM et al. Surveillance and analysis of avian influenza viruses, Australia. *Emerging Infectious Diseases*, 2010, 16:1896–1904. doi:10.3201/eid1612.100776 pmid:21122219
11. Takano R et al. Phylogenetic characterization of H5N1 avian influenza viruses isolated in Indonesia from 2003–2007. *Virology*, 2009, 390:13–21. doi:10.1016/j.virol.2009.04.024 pmid:19464724
12. Newcastle Disease – Papua New Guinea: (Sandaun) poultry. ProMed, 1 May 2013, archive no. 20130502.1685877 (<http://www.promedmail.org/?archiveid=20130502.1685877>, accessed 10 August 2013).
13. Horwood PF et al. Outbreak of chikungunya virus infection, Vanimo, Papua New Guinea. *Emerging Infectious Diseases*, 2013, 19:1535–1538. doi:10.3201/eid1909.130130 pmid:23965757
14. Olsen B et al. Global patterns of influenza A virus in wild birds. *Science*, 2006, 312:384–388. doi:10.1126/science.1122438 pmid:16627734
15. Roshier DA et al. Biogeographic models of gene flow in two waterfowl of the Australo-Papuan tropics. *Ecology and Evolution*, 2012, 2:2803–2814. doi:10.1002/ece3.393 pmid:23170215
16. Dingle H. The Australo-Papuan bird migration system: another consequence of Wallace's Line. *Emu*, 2004, 104:95–108. doi:10.1071/MU03026
17. Greenhill A et al. Improved laboratory capacity is required to respond better to future cholera outbreaks in Papua New Guinea. *Western Pacific Surveillance and Response Journal*, 2012, 3:30–32. doi:10.5365/wpsar.2011.2.4.016 pmid:23908909

Surveillance for arboviral zoonoses in New Zealand birds

Daniel Tompkins,^a Cheryl Johansen,^b Richard Jakob-Hoff,^c David Pulford,^d Isabel Castro^e and Graham Mackereth^f

Correspondence to Daniel Tompkins (e-mail: tompkinsd@landcareresearch.co.nz).

Introduction: Given the significant burden that emerging infectious diseases place on global economies and public health, the monitoring and mitigation of, and early response to, potential infectious diseases are of the highest priority. The objective of this study was to survey for known and other potential arboviral zoonoses in multiple bird species at four locations in New Zealand.

Methods: Common bird species were targeted for blood sampling during two southern hemisphere summers. Sera from each period ($n = 185$ and $n = 693$) were screened in an epitope blocking enzyme immunoassay for flavivirus antibody detection. In the first season, testing for antibodies to specific alphaviruses was conducted on samples with sufficient sera ($n = 22$). In the second season, blood clots ($n = 544$) were screened for viral presence by polymerase chain reaction (PCR) for alphaviral and flaviviral RNA, and virus isolation ($n = 146$) was conducted.

Results: Flavivirus antibodies were detected in 13 species, and one Australasian gannet (*Morus serrator*) from one site was positive for antibodies to Ross River virus. PCR tests and virus isolation were all negative.

Discussion: Evidence for flavivirus exposure in seabirds at Kaikoura Peninsula and Cape Kidnappers suggests that viruses isolated from seabirds and associated ticks in New Zealand in the late 1970s are still present. Evidence for flavivirus exposure in passerines at Kaikoura Peninsula, Cape Kidnappers and Mokoia Island is novel. The Ross River virus finding is also new and supports the hypothesis that migratory seabirds are an import pathway for such agents into New Zealand.

Emerging infectious diseases (EIDs; disease-causing agents that rapidly increase in host range, geographic range or prevalence) are a well-recognized threat to public health globally,¹ and the rate of disease emergence has risen since the middle of the 20th century.² Risk analysis indicates that emergence is driven by multiple factors including socioeconomic circumstances,^{2,3} climate and land-use changes,^{4,5} and pathogen pollution (the anthropogenic global movement of pathogens).⁶ Given the significant burden that EIDs place on global economies and public health,^{1,7} the monitoring and mitigation of, and early response to, potential infectious disease threats are of the highest priority.^{4,8} These global concerns are reflected in New Zealand with an increase in active surveillance for potential disease threats being advocated for the benefit of native wildlife, domestic stock and public health.^{9–15}

Four potential viral zoonoses associated with wildlife have previously been documented in New Zealand: three flaviviruses (Johnston Atoll virus,^{16,17} Saumarez

Reef virus and an unnamed Hughes group virus¹⁷) and one alphavirus (Whataroa virus¹⁸). The flaviviruses are all tick-borne viruses that have remained largely unstudied since their detection in the late 1970s. Johnston Atoll virus is closely related to the Quarantilla group of viruses, which have been isolated from symptomatic humans,¹⁶ and it has been hypothesized that humans may also be susceptible to infection with Johnston Atoll virus.^{16,19} Saumarez Reef virus is believed to have been responsible for febrile illness in meteorological workers on the Saumarez and Frederick reefs in Australia.²⁰ A closely related Hughes group virus, Soldado virus, has been implicated as a cause of human illness overseas.²¹ The Whataroa virus is a mosquito-borne alphavirus that belongs to the Sindbis virus subgroup that has had a known public health impact in several countries.²² Whataroa virus has been detected only in bird populations and two endemic mosquito species (*Culex pervigilans* and *Culiseta tonnoiri*) to date, around Whataroa township on New Zealand's South Island.¹⁸

^a Landcare Research, Dunedin, New Zealand.

^b Arbovirus Surveillance and Research Laboratory, School of Pathology and Laboratory Medicine, The University of Western Australia, Australia.

^c New Zealand Centre for Conservation Medicine, Auckland, New Zealand.

^d Investigation and Diagnostic Centres, Wallaceville, Upper Hutt, New Zealand.

^e Institute of Agriculture and Environment, Massey University, Palmerston North, New Zealand.

^f Institute of Environmental Science and Research, Porirua, New Zealand.

Submitted: 5 August 2013; Published: 6 November 2013

doi: 10.5365/wpsar.2013.4.3.002

The ecology and host-associations of all four viruses are poorly understood. In this study we conducted wildlife surveillance for these and other potential viral zoonoses at two locations where viruses were previously recorded (Kaikoura Peninsula and Cape Kidnappers; **Figure 1**) and two locations where occurrence was likely (Muriwai Beach for tick-borne viruses and Mokoia Island for mosquito-borne viruses). These locations are also potential import pathways for infectious agents into New Zealand; for example, migratory seabirds and their ticks may be able to transport infections such as West Nile virus into the country.²³ This potential import pathway has been discussed by various researchers globally,^{24–27} and the risk to New Zealand needs to be determined.


METHODS

Survey sites

The Kaikoura Peninsula, on the north-east coast of New Zealand's South Island, is where Saumarez Reef virus and the unidentified 'Hughes group' arbovirus were isolated from ticks associated with both the red-billed gull (*Larus novaehollandiae*) and white-fronted tern (*Sterna striata*) colonies in the 1970s and where the Hughes group virus was isolated from the blood of a red-billed gull.¹⁷ The presence of these viruses suggests a potential import pathway of migratory seabirds.²³ Red-billed gulls can move over 300 km after breeding, with some evidence of trans-oceanic straggling.²⁸ Large numbers of white-fronted terns migrate from New Zealand to Australia; the farthest recovery of a banded bird was 2970 km from Kaikoura to South Australia (**Figure 2**).²⁸

Cape Kidnappers, a peninsula on the east coast of New Zealand's North Island, has the country's largest mainland colony of the migratory Australasian gannet (*Morus serrator*). In the 1970s, Johnston Atoll virus was isolated from ticks associated with these gannets, in addition to the unidentified Hughes group arbovirus also isolated on the Kaikoura Peninsula.^{16,17} Most young Australasian gannets cross the Tasman Sea within three months of life,²⁸ remain in Australian waters until they are two to three years old (**Figure 2**), then return to their natal gannetries at three years of age as non-breeding or roosting birds – another potential import pathway.


Figure 1. Map of New Zealand indicating the four study locations



Muriwai Beach, on the west coast north of Auckland, has three potential import pathways. First, it is a second mainland colony site for migratory Australasian gannets; second, the site is close to major shipping ports and airports in the Auckland Region (both potential sites of entry of exotic vectors); and third, it is a popular tourist destination attracting thousands of overseas visitors each year. Being in the north of the country it also has close proximity to Australia and the Pacific islands (**Figure 2**).

Mokoia Island is a 1.35 km² island in the middle of Lake Rotorua in the centre of New Zealand's North Island. Infection of local bird populations by mosquito-borne avian malarial parasites have been documented here,²⁹ making it a potential site for mosquito-borne viral agents such as Whataroa virus. In addition, the migration of shining cuckoos (*Chrysococcyx lucidus*; a species that breeds on Mokoia Island) to the Bismarck (New Britain Island) and Solomon archipelagos and other Pacific Islands^{28,30} offers a potential route of agent incursion

Figure 2. Oceania regional map



(Figure 2). Mokoia Island is used for endangered bird translocations, representing a pathway for viral spread within the country.

Sampling

The common bird species present at each site were targeted for blood sampling during two southern hemisphere summers – January to March 2008 (all four sites) and November 2008 to February 2009 (Kaikoura Peninsula, Cape Kidnappers and Mokoia Island only). Tuis (*Prosthemadera novaeseelandiae*), North Island robins (*Petroica longipes*), North Island saddlebacks (*Philesturnus rufusater*) and other passerines were caught using mist nets, banded with a numbered metal band (if no band already present) and had a peripheral blood sample collected from the brachial vein. The vein was punctured using a sterile 25–27 g needle (depending on bird size), and blood (no more than 1% body weight) was collected into capillary tubes.

Hand nets were used to catch red-billed gulls and white-fronted terns, and shepherd's crooks were used to catch Australasian gannets. Little blue penguins (*Eudyptula minor*) were taken by hand from burrows as were gulls and terns from nests. Wekas (*Gallirallus australis*) were caught in baited cage-traps, and

New Zealand scaup (*Aythya novaeseelandiae*) were caught in mist nets on the shore of Lake Rotorua (in which Mokoia Island lies). Once banded with a numbered metal band (if no band already present), a peripheral blood sample was collected. Gannets, penguins, gulls, terns, scaups and wekas had up to 1.0 ml blood drawn by syringe with a sterile 25 g needle from the metatarsal vein. Gulls and juvenile terns had their brachial vein punctured using a sterile 25 g needle with up to 0.5 ml blood collected into capillary tubes.

Diagnostic testing

Serum samples (collected from $n = 185$ and $n = 693$ individuals during the first and second field seasons respectively) were screened using an flavivirus epitope-blocking enzyme-linked immunosorbent assay described elsewhere^{31,32} with the exception that virus-inactivated cell culture lysates were used to coat U-bottom 96-well plates before addition of test samples.³³ Briefly, after washing excess antigen and blocking, sera were added to the 96-well plates in duplicate before the addition of the flavivirus group-reactive monoclonal antibody 3H6 (JCU Tropical Biotechnology Pty Ltd, Townsville, Australia). Binding of the monoclonal antibody was detected following the addition of horseradish peroxidase-conjugated goat anti-mouse antibody and subsequent visualization of enzymatic activity in substrate buffer.

Optical densities were measured and percentage inhibition of the monoclonal antibody by test sera was calculated using negative control sera as the reference. For samples with sufficient sera, those with 30% or greater inhibition were re-tested against 3H6 as well as specific monoclonal antibodies 10C6 (JCU Tropical Biotechnology Pty. Ltd) for Murray Valley encephalitis virus and 3.1112G (Discipline of Microbiology and Immunology, The University of Western Australia, Perth, Australia) for Kunjin virus (both flaviviral agents of incursion concern from Australia¹⁴). Samples with 50% or greater inhibition on at least one 3H6 test were considered positive for flavivirus antibodies. This criterion was validated as robust in the 50 samples that were re-tested; while some samples up to 40% did not confirm at re-testing, all samples over 40% did.

Testing for antibodies to specific alphaviruses (Ross River virus, Barmah Forest virus and Sindbis virus; arboviral agents of incursion concern from Australia^{14,34}) was also carried out on first field season samples with sufficient remaining sera ($n = 22$) using serum neutralization assays as described elsewhere³⁵ except that Vero cells were used in place of baby hamster kidney cells. In short, sera were serially diluted in 96-well tissue culture plates and incubated for five days with approximately 50 tissue culture infectious doses of virus and Vero cells. Each well was examined microscopically for cytopathic effect (CPE), and neutralization titres were expressed as the reciprocal of the highest serum dilution where CPE did not occur. Samples with two repeat neutralization titres of at least 40 were considered positive.

Blood clots collected during the second field season (from $n = 544$ individuals) were screened for viral presence using flavivirus and alphavirus group-specific reverse transcription–PCR tests. Blood clots collected after removal of serum were frozen at -70°C within four hours of collection. They were then homogenized in sterile virus transport media and the debris pelleted by microcentrifugation. The collected supernatant was extracted directly using the Zymo Viral RNA kit (Zymo Research Corporation, Irvine, CA, USA) and resuspended in ddH₂O. For flavivirus testing, we employed the flavivirus nsp5 PCR that uses mFU1 and cFD2 published primers.^{36,37} Generic alphavirus PCR was conducted using nsp4 AL-EF and AL-ER primers.³⁸ Both PCR tests were conducted using

Invitrogen Superscript III Platinum Taq Sybr Green one-step qRT–PCR master mix (Life Technologies, Carlsbad, CA, USA) in single-tube reactions. The detection of positive reactions was determined by melt curve analysis of the PCR product followed by gel electrophoresis and DNA sequencing of PCR amplicons.

Virus isolation was performed on clots collected from 146 individuals during the second field season. Supernatants from homogenized clots were inoculated into VeroE6 cell cultures³⁹ for two passes of five days and monitored for evidence of CPE using a light microscope.

RESULTS

Serology

In the first field season, antibodies to flavivirus were detected in serum samples from a red-billed gull at Kaikoura Peninsula and a North Island saddleback at Mokoia Island (**Table 1**). In the second field season, a relatively high prevalence of antibodies to flavivirus was observed in serum samples from white-fronted terns at Kaikoura Peninsula (**Table 1**). Flavivirus antibodies were also detected at this time in red-billed gulls and passerines at this location; in little blue penguins and passerines at Cape Kidnappers; and in passerines, wekas and New Zealand scaups at Mokoia Island (**Table 1**). None of the 50 repeat-tested samples were specifically positive for either Murray Valley encephalitis virus or Kunjin virus.

Of the 22 first field season samples also tested for antibodies to specific alphaviruses (**Table 2**), one Australasian gannet from Muriwai Beach was positive for antibodies to Ross River virus with two repeat neutralization titres of 80.

PCR and virus isolation

In the second field season, the 544 blood clots (from Kaikoura Peninsula, Cape Kidnappers and Mokoia Island) screened on alphavirus and flavivirus generic PCR tests were all negative (**Table 3**). The 146 clots subject to viral isolation were also negative (**Table 3**); no CPE was observed in any of the cultures after two passages of virus isolation in VeroE6 cells, and no flavivirus PCR products were amplified with RNA extracted from these cell cultures.

Table 1. Confirmed flavivirus antibody-positive serum samples collected from birds in both the first (2007/08 southern hemisphere summer) and second (2008/09 southern hemisphere summer) field seasons

Common name	Latin name	Number of individuals positive for flavivirus neutralizing antibodies/Total number screened								Total
		Cape Kidnappers		Muriwai Beach		Kaikoura Peninsula		Mokoia Island		
		2007/08	2008/09	2007/08	2008/09	2007/08	2008/09	2007/08	2008/09	
Australian magpie	Gymnorhina tibicen	–	0/1	–	–	–	–	–	–	0/1
Australasian gannet	Gallirallus australis	0/35	0/131	0/57	–	–	–	–	–	0/223
Chaffinch	Fringilla coelebs	-	0/1	–	–	–	0/7	–	–	0/8
Cirl bunting	Emberiza cirlus	-	-	–	–	–	1/1	–	–	1/1
Common starling	Sturnus vulgaris	0/1	0/1	–	–	–	0/3	–	–	0/5
Dunnock	Prunella modularis	0/1	0/5	–	–	–	0/4	–	–	0/10
Eurasian blackbird	Turdus merula	0/2	0/3	–	–	–	1/14	0/1	2/10	3/30
European goldfinch	Carduelis carduelis	-	0/2	–	–	–	0/4	–	–	0/6
Greenfinch	Carduelis chloris	0/3	0/3	–	–	–	1/5	–	–	1/11
House sparrow	Passer domesticus	0/6	0/34	–	–	–	0/11	–	–	0/51
Little blue penguin	Eudyptula minor	–	2/17	–	–	0/7	0/10	–	–	2/34
New Zealand scaup	Aythya novaeseelandiae	–	–	–	–	–	–	–	1/12	1/12
North Island robin	Petroica longipes	–	–	–	–	–	–	0/15	1/38	1/53
North Island saddleback	Philesturnus rufusater	–	–	–	–	–	–	1/38	0/77	2/115
Red-billed gull	Larus novaehollandiae	–	0/18	–	–	1/15	6/104	–	–	7/137
Silvereye	Zosterops lateralis	0/3	0/11	–	–	–	–	–	–	0/14
Song thrush	Turdus philomelos	–	1/3	–	–	–	0/7	–	0/3	1/13
Tui	Prosthemadera novaeseelandiae	–	–	–	–	–	–	0/1	2/28	2/29
Weka	Gallirallus australis	–	–	–	–	–	–	–	1/8	1/8
Welcome swallow	Hirundo neoxena	–	0/3	–	–	–	–	–	–	0/3
White-fronted tern	Sterna striata	–	–	–	–	–	33/102	–	–	33/102
Yellowhammer	Emberiza citrinella	–	1/4	–	–	–	1/8	–	–	2/12
Site totals		4/288		0/57		44/302		8/231		

DISCUSSION

The four sites surveyed for viral agents in birds were selected on the basis of previous documentation of potential zoonoses (in seabirds and their associated ticks) and/or the presence of potential import pathways. Our results indicate that these selection criteria were relevant. Evidence suggests the continued presence of previously isolated seabird flaviviruses, the presence of novel avian flaviviral agents and exposure of a migratory species to an alphavirus of incursion concern from Australia. This last result, serological evidence for antibodies to Ross River virus (the most common mosquito-borne pathogen causing human disease in Australia³⁴) in an Australasian gannet at Muriwai Beach, is a novel finding of particular relevance to public health.

Although the standard positive criterion for the flavivirus serology conducted is to achieve inhibition of 3H6 on repeat testing, we were frequently unable to obtain sufficient serum for a repeat (particularly from smaller birds). To maximize the utility of our surveys, and prevent biasing against smaller species in our findings, we instead used a criterion of 50% or greater inhibition on at least one test. Although this criterion was validated as robust in the 50 samples that were re-tested (while some samples up to 40% did not confirm at re-testing, all samples over 40% did), our inability to conduct repeat testing on all samples means that cases of just one or two positive results should be interpreted with caution and require follow-up sampling to confirm the evidence for flavivirus infection. In spite of this proviso, we have obtained two strong lines of evidence for such infection.

Table 2. **Confirmed alphavirus antibody-positive serum samples collected from birds in the first field season (2007/08 southern hemisphere summer) for antibodies to specific alphaviruses***

Common name	Location	Number positive	Number negative
Australasian gannet	Cape Kidnappers	0	4
Australasian gannet	Muriwai Beach	1 (RRV)	12
Little blue penguin	Kaikoura Peninsula	0	1
Red-billed gull	Kaikoura Peninsula	0	3
North Island saddleback	Mokoia Island	0	1

* Specific alphaviruses – Ross River virus (RRV), Barmah Forest virus, Sindbis virus. See Table 1 for species Latin names.

Table 3. **Blood clots collected in the second field season (2008/09 southern hemisphere summer) subjected to alphaviral and flaviviral PCR assays and virus isolation***

Common name	Number of individuals screened for alphaviruses/flaviviruses			Number of individuals screened by virus isolation	
	Cape Kidnappers	Mokoia Island	Kaikoura Peninsula	Cape Kidnappers	Mokoia Island
Australian magpie	–	–	–	1	–
Australasian gannet	54/54	–	–	22	–
Chaffinch	–	–	11/12	1	–
Cirl bunting	–	–	0/1	–	–
Common starling	–	–	1/3	1	–
Dunnock	3/3	–	8/8	4	–
Eurasian blackbird	–	2/2	17/21	2	1
European goldfinch	–	–	5/6	1	–
Greenfinch	–	–	2/7	1	–
House sparrow	9/9	–	34/35	24	–
Little blue penguin	–	–	11/11	17	–
North Island robin	–	30/46	–	–	12
North Island saddleback	–	54/54	–	–	20
Red-billed gull	1/1	–	119/119	11	–
Silvereye	10/10	–	–	12	–
Song thrush	2/2	1/1	8/10	3	1
Tui	–	6/6	–	–	3
Weka	–	3/3	–	–	3
Welcome swallow	–	–	–	3	–
White-fronted tern	–	–	0/103	–	–
Yellowhammer	2/2	–	13/15	3	–
Site totals	81/81	96/112	228/351	106	40

* All tests were negative. See Table 1 for species Latin names.

First, serology results from Kaikoura Peninsula suggest that previously isolated flaviviruses from red-billed gulls (the unidentified Hughes group arbovirus) and ticks associated with both red-billed gulls and white-fronted terns (Saumarez Reef virus and the unidentified Hughes group arbovirus) are still present at this site. Targeted sampling at different times of year may be required for successful viral isolation to verify agent identity. With

specific tests for flaviviral agents of incursion concern being negative, the flaviviral reactivity detected in little blue penguins at Cape Kidnappers similarly suggests that the viruses previously isolated from ticks associated with Australasian gannets at this site (Johnston Atoll virus and the unidentified Hughes group arbovirus) may also still be present. However, successful viral isolation is again necessary to verify this.

Second, serological evidence for flavivirus exposure in passerines is novel with no prior evidence for such agents being present in such hosts. Targeted sampling at different times of year may once again be required for successful viral isolation to identify the agents present and inform whether this represents a past incursion via a migratory species such as the shining cuckoo. Since human flaviviral infection is as yet unknown in New Zealand,⁹ these agents are most likely not a risk to public health.

CONCLUSIONS

The key conclusion that can be drawn from both the results discussed above and previous work is that migratory birds represent a possible import pathway for potential zoonotic agents into New Zealand. Both the past and current evidence for Saumarez Reef virus and Johnston Atoll virus support the hypothesis that this pathway has historically operated to bring such agents into the country. Although birds may not be currently carrying viral particles back into New Zealand, the evidence for Australasian gannet exposure to Ross River virus indicates that incursion from Australia by such a mechanism may be possible. Since the native *Aedes notoscriptus* and *Culex pervigilans* and the introduced *Aedes camptorhynchus*, *Aedes australis* and *Culex quinquefasciatus* mosquitoes are all potentially competent vectors of Ross River virus,^{14,40} such incursion could lead to ongoing transmission within the country. With this agent being of public health concern, more thorough surveillance should be carried out at Muriwai Beach to confirm its current status.

Conflicts of interest

None declared.

Funding

This work was funded by the New Zealand Foundation for Research Science and Technology (now Ministry of Business, Innovation and Employment).

Acknowledgements

Thanks to Lindsay Rowe and Jim Mills for field assistance at Kaikoura Peninsula, to Raewyn Edmonds and Te Rūnanga o Kaikoura for iwi approval to work at

Kaikoura Peninsula, to Mick Unahi and Ngāti Hawea for iwi approval to work at Cape Kidnappers, to Malcolm Paterson and Ngāti Whatua o Kaipara for iwi approval to work at Muriwai Regional Park, and to Bill Kingi and the Mokoia Island Trust for iwi approval to work on Mokoia Island. Thanks also to Dean Clarke, Morgan Coleman, Keven Drew, Steph Hicks, Pete Lei, Adrian Monks, Maria Barclay, Lauren Best, Kirsten Derry, Mel Farrant, John Potter, Stephanie Shaw, Ellen Schoener, Cleland Wallace, Stefanie Ismar and Katja Geschke for field assistance. Further thanks to Della Orr for help with virology test development, Megan Dymond and Jianing Wang for contributions to PCR test development and Cheryl Johansen for serological testing. This work was conducted under New Zealand Department of Conservation (DOC) Global Concession CA-5160-OTH; DOC Research and Collection Permits NM-22225-RES, ECHB-22299-FAU, AK-22099-FAU, BP-22190-RES, NM-23980-RES, ECHB-24005-FAU and BP-23988-RES; Landcare Research Animal Ethics Authority 07/12/01; New Zealand National Bird Banding Scheme Institutional Permit to Band Birds No. 2007/83.

References

- Morens DM, Folkers GK, Fauci AS. The challenge of emerging and re-emerging infectious diseases. *Nature*, 2004, 430:242–249. doi:10.1038/nature02759 pmid:15241422
- Jones KE et al. Global trends in emerging infectious diseases. *Nature*, 2008, 451:990–993. doi:10.1038/nature06536 pmid:18288193
- Weiss RA, McMichael AJ. Social and environmental risk factors in the emergence of infectious diseases. *Nature Medicine*, 2004, 10 Suppl:S70–76. doi:10.1038/nm1150 pmid:15577934
- Patz JA et al.; Working Group on Land Use Change and Disease Emergence. Unhealthy landscapes: Policy recommendations on land use change and infectious disease emergence. *Environmental Health Perspectives*, 2004, 112:1092–1098. doi:10.1289/ehp.6877 pmid:15238283
- Patz JA et al. Impact of regional climate change on human health. *Nature*, 2005, 438:310–317. doi:10.1038/nature04188 pmid:16292302
- Cunningham AA, Daszak P, Rodriguez JP. Pathogen pollution: defining a parasitological threat to biodiversity conservation. *Journal of Parasitology Archives*, 2003, 89:S78–S83.
- Meslin FX, Stöhr K, Heymann D. Public health implications of emerging zoonoses. *Revue Scientifique et Technique (International Office of Epizootics)*, 2000, 19:310–317. pmid:11189723
- King DA et al. Epidemiology. Infectious diseases: preparing for the future. *Science*, 2006, 313:1392–1393. doi:10.1126/science.1129134 pmid:16959992
- Crump JA, Murdoch DR, Baker MG. Emerging infectious diseases in an island ecosystem: the New Zealand perspective. *Emerging Infectious Diseases*, 2001, 7:767–772. pmid:11747690
- Alley MR. Avian wildlife diseases in New Zealand: current issues and achievements. *New Zealand Veterinary Journal*, 2002, 50 Suppl:118–120. doi:10.1080/00480169.2002.36287 pmid:16032257

11. Tompkins DM, Poulin R. Parasites and biological invasions. In: Allen RB, Lee WG, eds. *Biological invasions in New Zealand*. Ecological Studies 186. Berlin, Springer, 2006, 67–84.
12. Derraik JGB, Slaney D. Anthropogenic environmental change, mosquito-borne diseases and human health in New Zealand. *EcoHealth*, 2007, 4:72–81. doi:10.1007/s10393-006-0080-2
13. French NP, Gemmell NJ, Buddle BM. Advances in biosecurity to 2010 and beyond: towards integrated detection, analysis and response to exotic pest invasions. *New Zealand Veterinary Journal*, 2007, 55:255–263. doi:10.1080/00480169.2007.36779 pmid:18059642
14. Mackereth G et al. *Vectors and vector borne diseases: ecological research and surveillance development in New Zealand. Risk assessment*. Wellington, MAF(BNZ), 2007.
15. Derraik JGB, Calisher CH. Is New Zealand prepared to deal with arboviral diseases? *Australian and New Zealand Journal of Public Health*, 2004, 28:27–31. doi:10.1111/j.1467-842X.2004.tb00628.x pmid:15108743
16. Austin FJ. Johnston Atoll virus (Quarantilla group) from *Ornithodoros capensis* (Ixodoidea: Argasidae) infesting a gannet colony in New Zealand. *The American Journal of Tropical Medicine and Hygiene*, 1978, 27:1045–1048. pmid:717630
17. Austin FJ. Ticks as arbovirus vectors in New Zealand. *New Zealand Entomologist*, 1984, 8:105–106. doi:10.1080/00779962.1984.9722481
18. Tompkins DM et al. Whataroa virus four decades on – Emerging, persisting, or fading out? *Journal of the Royal Society of New Zealand*, 2010, 40:1–9. doi:10.1080/03036751003641701
19. Clifford CM et al. Identification and comparison of two viruses isolated from ticks of the genus *Ornithodoros*. *The American Journal of Tropical Medicine and Hygiene*, 1968, 17:881–885. pmid:4973055
20. St George TD et al. The isolation of Saumarez Reef virus, a new flavivirus, from bird ticks *Ornithodoros capensis* and *Ixodes euryptidis* in Australia. *The Australian Journal of Experimental Biology and Medical Science*, 1977, 55:493–499. doi:10.1038/icb.1977.49 pmid:75000
21. Chastel C, Bailly-Choumara H, Le Lay G. [Natural pathogenicity for man of an antigenic variant of Soldado virus from Morocco (author's transl)]. *Bulletin de la Société de Pathologie Exotique*, 1981, 74:499–505. pmid:6274527
22. Saleh SM et al. Antigenic and genetic typing of Whataroa viruses in Australia. *The American Journal of Tropical Medicine and Hygiene*, 2004, 71:262–267. pmid:15381803
23. Heath ACG. A review of the origins and zoogeography of tick-borne disease in New Zealand. *Tuatara*, 1987, 29:19–29.
24. Olsen B et al. Transhemispheric exchange of Lyme disease spirochetes by seabirds. *Journal of Clinical Microbiology*, 1995, 33:3270–3274. pmid:8586715
25. Rappole JH, Derrickson SR, Hubálek Z. Migratory birds and spread of West Nile virus in the Western Hemisphere. *Emerging Infectious Diseases*, 2000, 6:319–328. doi:10.3201/eid0604.000401 pmid:10905964
26. Peterson AT, Vieglais DA, Andreasen JK. Migratory birds modeled as critical transport agents for West Nile Virus in North America. *Vector Borne and Zoonotic Diseases (Larchmont, N.Y.)*, 2003, 3:27–37. doi:10.1089/153036603765627433 pmid:12804378
27. Reed KD et al. Birds, migration and emerging zoonoses: west nile virus, lyme disease, influenza A and enteropathogens. *Clinical Medicine & Research*, 2003, 1:5–12. doi:10.3121/cmr.1.1.5 pmid:15931279
28. Williams M et al. *Migrations and movements of birds to New Zealand and surrounding seas*. Wellington, Department of Conservation Science and Research Unit Report to MAF, Biosecurity Authority, 2004.
29. Castro I et al. Presence and seasonal prevalence of *Plasmodium* spp. in a rare endemic New Zealand passerine (tieke or Saddleback, *Philesturnus carunculatus*). *Journal of Wildlife Diseases*, 2011, 47:860–867. doi:10.7589/0090-3558-47.4.860 pmid:22102656
30. Heather B, Robertson H. *The field guide to the birds of New Zealand*. Auckland, Viking, 2005.
31. Hall RA et al. Immunodominant epitopes on the NS1 protein of MVE and KUN viruses serve as targets for a blocking ELISA to detect virus-specific antibodies in sentinel animal serum. *Journal of Virological Methods*, 1995, 51:201–210. doi:10.1016/0166-0934(94)00105-P pmid:7738140
32. Broom AK et al. Investigation of the southern limits of Murray Valley encephalitis activity in Western Australia during the 2000 wet season. *Vector Borne and Zoonotic Diseases (Larchmont, N.Y.)*, 2002, 2:87–95. doi:10.1089/153036602321131887 pmid:12653302
33. Spicer PE et al. Antibodies to Japanese encephalitis virus in human sera collected from Irian Jaya. Follow-up of a previously reported case of Japanese encephalitis in that region. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 1999, 93:511–514. doi:10.1016/S0035-9203(99)90353-X pmid:10696406
34. Tompkins DM, Slaney D. Exploring the potential for Ross River virus emergence in New Zealand. *Vector Borne and Zoonotic Diseases (Larchmont, N.Y.)*. In press.
35. Johansen CA et al. Prevalence of neutralising antibodies to Barmah Forest, Sindbis and Trubanaman viruses in animals and humans in the south-west of Western Australia. *Australian Journal of Zoology*, 2005, 53:51–58. doi:10.1071/Z003042
36. Kuno G. Universal diagnostic RT-PCR protocol for arboviruses. *Journal of Virological Methods*, 1998, 72:27–41. doi:10.1016/S0166-0934(98)00003-2 pmid:9672130
37. Chao DY, Davis BS, Chang GJ. Development of multiplex real-time reverse transcriptase PCR assays for detecting eight medically important flaviviruses in mosquitoes. *Journal of Clinical Microbiology*, 2007, 45:584–589. doi:10.1128/JCM.00842-06 pmid:17108075
38. Sánchez-Seco MP et al. A generic nested-RT-PCR followed by sequencing for detection and identification of members of the alphavirus genus. *Journal of Virological Methods*, 2001, 95:153–161. doi:10.1016/S0166-0934(01)00306-8 pmid:11377722
39. Earley EM, Johnson KM. The lineage of Vero, Vero 76 and its clone C1008 in the United States In: Earley EM, Johnson KM, *Vero cells: origin, properties and biomedical applications*. Tokyo, Chiba University, 26–29, 1988.
40. Kramer LD et al. Vector competence of New Zealand mosquitoes for selected arboviruses. *The American Journal of Tropical Medicine and Hygiene*, 2011, 85:182–189. doi:10.4269/ajtmh.2011.11-0078 pmid:21734146

Human infections with avian influenza A(H7N9): preliminary assessments of the age and sex distribution

Viroj Wiwanitkit^a

The recent report on human infections with avian influenza A(H7N9) is very interesting.¹

The age distribution of the patients was studied, and Arima et al. mentioned that “it seems unlikely that elderly men are being overly selected.”¹ It is true that the virus can attack any age group. The factors that determine vulnerability to infection in each age group include (1) immunity to infection, (2) exposure to the disease, (3) availability of medical care, and (4) ability of medical personnel to diagnose the illness.

The elderly are more prone to infections due to their weaker health status as compared to the young. In addition, the high number of infections among the elderly might imply that they have no previous immunity to the infection, indicating that avian influenza A(H7N9) is a new infection for the Chinese in our generation (compared to swine flu in which there was evidence of cross-protective immunity among the elderly

that might relate to the low number of cases among that group²).

Conflicts of interest

None declared.

Funding

None.

References

1. Arima Y, Vong S; World Health Organization Outbreak Response Team. Human infections with avian influenza A(H7N9) virus in China: preliminary assessments of the age and sex distribution. *Western Pacific Surveillance and Response Journal*, 2013, 4(2): 1–3. doi:10.5365/wpsar.2013.4.2.005 pmid:24015363
2. De Marco MA et al. Evidence of cross-reactive immunity to 2009 pandemic influenza A virus in workers seropositive to swine H1N1 influenza viruses circulating in Italy. *PLoS ONE*, 2013, 8:e57576. doi:10.1371/journal.pone.0057576 pmid:23469029

^a Hainan Medical University, People's Republic of China (e-mail: vviroj@yahoo.com).
Submitted: 13 September 2013; Published: 31 October 2013
doi: 10.5365/wpsar.2013.4.3.006

Author Response

Yuzo Arima,^a Rongqiang Zu,^a Manoj Murhekar,^a Sirenda Vong^b and Tomoe Shimada^a

Correspondence to Yuzo Arima (e-mail: arimay@wpro.who.int).

We thank Dr Viroj Wiwanitkit for his comments on our preliminary assessment of the age and sex distribution of the human cases with avian influenza A(H7N9) virus infection. To clarify, we posed three scenarios which could possibly explain the preponderance of cases among elderly men reported through China's surveillance system: (1) differential exposure due to gender-associated practices and norms, e.g. more high-risk behaviours among elderly men; (2) differential clinical course post-exposure/

infection, e.g. given similar exposures, elderly men have a more severe outcome relative to other age-gender groups; and (3) differential health care-seeking/access behaviour favouring selection of elderly men, e.g. elderly men accessing health care more than other age-gender groups.¹ There may be more than one of these possibilities in operation, and this initial assessment was intended to pose the question to public health practitioners and researchers and to encourage further study into the causes for the distribution observed for

^a Division of Health Security and Emergencies, World Health Organization Regional Office for the Western Pacific, Manila, Philippines.

^b Emerging Disease Surveillance and Response, World Health Organization China Office, Beijing, China.

Submitted: 14 October 2013; Published: 31 October 2013

doi: 10.5365/wpsar.2013.4.4.003

this apparently emerging disease.² Our statement that it appeared unlikely that elderly men were being overly selected was addressing this third possibility.

While we agree with the four possibilities listed by Dr Wiwanitkit “that determine vulnerability to infection”, we also believe that vulnerability to severe outcomes (i.e. scenario 2, differential outcomes given an infection) is important when assessing surveillance information given that reported surveillance data are often a function of severity. For example, during the 2009 H1N1 pandemic, while infection rates were lower in the elderly (attributed to likely cross-protection from previous H1N1 infection among the elderly survivors) relative to seasonal influenza, once infected, the elderly were still at higher risk of serious complications.³ In addition, a recent serological study in China found that, while no seropositivity for antibodies specific for H7N9 virus were detected among >1000 individuals among the general population, greater than 6% of the 396 poultry workers tested were positive, indicating that subclinical or non-severe infections are possible.⁴

We agree with Dr Wiwanitkit's statement that “the virus can attack any age group”. As we reported (age range 4–87 years) and as reported later in August 2013, China's routine influenza-like illness surveillance detected from outpatient visits six avian influenza A(H7N9) cases that skewed towards a younger profile.⁵ Of these, four had complications and were hospitalized. Notably, the non-hospitalized cases were aged two and four years, while those hospitalized were older. This adds to the biological possibility that, once infected with avian influenza A(H7N9), the elderly may suffer more severe outcomes relative to their younger cohort. While acknowledging the wide age range for infection, the distribution of the avian influenza A(H7N9) cases continues to tend towards the elderly (more than half of cases are 60 years or older as of late September 2013, $n = 135$); this distribution remains strikingly different from that of avian influenza A(H5N1) and requires further investigation. As we noted regarding seasonal influenza infections, the elderly are generally more prone to suffer from severe clinical manifestation of influenza virus infection,^{3,6,7} and this may be the case for avian influenza A(H7N9).

Lastly, we appreciate Dr Wiwanitkit's comment regarding the possibility of the absence of immunity to the avian influenza A(H7N9) virus among the elderly, hence the true novel nature of the avian influenza A(H7N9)

virus in humans causing infection in all ages rather than a detection and/or reporting artefact. We agree with this statement that is supported through extensive phylogenetic and virological analyses,⁸ the absence of pre-existing immunity to avian influenza A(H7N9) among high-risk groups before 2013⁹ and the lack of cross-reactive immunity in tested patients previously vaccinated against seasonal influenza viruses.¹⁰

As the winter influenza season in the northern hemisphere approaches with the potential for additional cases of avian influenza A(H7N9), it is imperative that investigations continue with regards to the observed skewed age and sex distributions.

Conflicts of interest

None declared.

Funding

None.

References:

1. Arima Y et al. Human infections with avian influenza A(H7N9) virus in China: preliminary assessments of the age and sex distribution. *Western Pacific Surveillance and Response*, 2013, 4(2):1–3. doi:10.5365/wpsar.2013.4.2.005 pmid:24015363
2. Cowling BJ et al. Comparative epidemiology of human infections with avian influenza A H7N9 and H5N1 viruses in China: a population-based study of laboratory-confirmed cases. *Lancet*, 2013, 382:129–137. doi:10.1016/S0140-6736(13)61171-X pmid:23803488
3. *Updated CDC Estimates of 2009 H1N1 Influenza Cases, Hospitalizations and Deaths in the United States, April 2009 – April 10, 2010*. Atlanta, Centers for Disease Control and Prevention, 2010 (http://www.cdc.gov/h1n1flu/estimates_2009_h1n1.htm, accessed 20 September 2013).
4. Yang S et al. Avian-origin influenza A(H7N9) infection in influenza A(H7N9)-affected areas of China: a serological study. *Journal of Infectious Diseases*, 2013, Sep 18. pmid:23935201
5. Xu C et al. Monitoring avian influenza A(H7N9) virus through national influenza-like illness surveillance, China. *China Emerging Infectious Diseases*, 2013, 19:1289–1292. pmid:23879887
6. Molinari NA et al. The annual impact of seasonal influenza in the US: measuring disease burden and costs. *Vaccine*, 2007, 25:5086–5096. doi:10.1016/j.vaccine.2007.03.046 pmid:17544181
7. Eshima N et al. Sex- and age-related differences in morbidity rates of 2009 pandemic influenza A H1N1 virus of swine origin in Japan. *PLoS ONE*, 2011, 6:e19409. doi:10.1371/journal.pone.0019409 pmid:21559366
8. Liu D et al. Origin and diversity of novel avian influenza A H7N9 viruses causing human infection: phylogenetic, structural, and coalescent analyses. *Lancet*, 2013, 381:1926–1932. doi:10.1016/S0140-6736(13)60938-1 pmid:23643111
9. Bai T, Zhou J, Shu Y. Serologic study for influenza A (H7N9) among high-risk groups in China. *The New England Journal of Medicine*, 2013, 368:2339–2340. doi:10.1056/NEJMc1305865 pmid:23718151
10. Zhou J et al. Biological features of novel avian influenza A (H7N9) virus. *Nature*, 2013, 499:500–503. doi:10.1038/nature12379 pmid:23823727

Western Pacific Surveillance and Response

Instructions to Authors

ABOUT WPSAR

The aims of WPSAR are:

1. to provide an open access journal to publish articles on the surveillance of and response to public health events and emergencies in the WHO Western Pacific Region and in areas with relevance to the Western Pacific Region; and
2. to build capacity in communicating epidemiological and operational research within the WHO Western Pacific Region.

Our objectives are:

1. to provide a platform for people working in surveillance and response in the Western Pacific Region to share their scientific and operational findings;
2. to publish a broad range of articles not limited to conventional research articles:
 - to disseminate short reports on outbreak investigations;
 - to publish analyses of surveillance data on communicable diseases;
 - to encourage the publication of evaluations of new and existing surveillance systems;
 - to promote the use of risk assessment for public health by facilitating risk assessment articles;
 - to support preparedness and response to public health events and emergencies through the dissemination of lessons learnt from such events; and
3. to build capacity in communicating epidemiological and operational findings in the Western Pacific Region through pre-submission assistance.

Scope

WPSAR covers all activities related to the surveillance of and response to public health events and emergencies, with a focus on topics that are relevant to the Western Pacific Region. Public health events may be acute or ongoing and can fall under any of the following areas: communicable diseases, natural disasters, food safety, bioterrorism, and chemical and radiological events. Other events and topics may also be considered. Response activities include those for acute events, e.g. responding to natural disasters, or for response to cases or epidemics of disease.

Why publish in WPSAR?

WPSAR is not limited to conventional research. It publishes a broad range of articles, including short outbreak investigation reports, lessons from the field, analyses of surveillance data, evaluations of surveillance systems and risk assessments for public health events. There are limited opportunities to publish these types of articles in other journals. We also accept the more traditional original research, perspectives and case reports/case series articles.

WPSAR is an open access journal, meaning it is free of charge for both readers and authors. It is also a continuous publication, which means articles are published as soon as they have completed the review and editing process.

WPSAR accepts all articles that fit the scope of the journal and that meet the minimum publication standards. We are especially interested in field epidemiology and operational research.

WPSAR also aims to build capacity in scientific writing and encourages submissions from authors with little or no experience in publishing in peer-reviewed journals. The Coordinating Editor often works with new authors on their submissions to ensure that articles fit the scope of WPSAR and meet the minimum standards for publication.

INSTRUCTIONS TO AUTHORS FOR ARTICLE WRITING AND SUBMISSION

WPSAR follows the guidelines of the *Uniform Requirements for Articles Submitted to Biomedical Journals by the International Committee for Medical Journal Editors (ICMJE)*.

Formatting guidelines

Please submit your article in a Microsoft® Office Word file or a compatible file in English. Double-spaced, 12-point Arial font should be used to format your article. Please remove all automatic formatting including automatic numbering and referencing before submitting.

The format of the article will depend on the article type. Please see below for specific instructions per article type.

Outbreak Investigation Report

A short article describing a field or outbreak investigation including how it was detected, investigated and controlled. Rapid risk assessments undertaken during these investigations are also encouraged. These articles may be considered for rapid publication.

- Structured article with an abstract of ≤ 250 words and sections for introduction, methods, results and discussion
- Structured abstract with sections for objective, methods, results and discussion
- Word limit: ≤ 1500 words
- ≤ 15 references
- ≤ 2 figures/graphs/pictures

More comprehensive investigations can be submitted as Original Research.

Surveillance Report

A summary and interpretation of surveillance data over a given period of time. A description of the surveillance system and the limitations of the data collected must be included.

- Unstructured abstract of ≤ 250 words
- Word limit: ≤ 2000 words
- ≤ 15 references
- ≤ 10 figures/graphs/pictures

Surveillance System Implementation/Evaluation

An article describing the implementation of a new surveillance system or an evaluation of an existing surveillance system used to detect public health events.

- Unstructured abstract of ≤ 250 words
- Word limit: ≤ 2000 words
- ≤ 15 references
- ≤ 3 figures/graphs/pictures

Risk Assessments

An article detailing a risk assessment of a public health threat or event.

- Structured article with an abstract ≤ 250 words and sections for introduction (including risk question/s), risk assessment methodology, results, discussion and recommendations
- Structured abstract with objectives, method, results and discussion
- The results should include an assessment and/or characterization of the hazard, exposure and context, as well as the level of risk or risk characterization. The limitations must also be included. Risk management may be included in the discussion.
- Word limit: ≤ 3000 words
- ≤ 30 references
- ≤ 3 figures/graphs/pictures

Original Research

Original research articles may include epidemiological studies including outbreak investigations.

- Structured article with an abstract of ≤ 250 words and sections for introduction, methods, results and discussion
- Structured abstract with objective, methods, results and discussion
- Word limit: ≤ 3000 words
- ≤ 40 references
- ≤ 5 figures/graphs/pictures

Lessons from the Field

An article describing a problem faced in field epidemiology or during a public health event and the experience in trying to overcome the problem.

- Structured article with an abstract ≤ 250 words and sections for problem, context, action, lesson(s) learnt or outcome and discussion
- Structured abstract with the headings of problem, context, action, lesson(s) learnt and discussion
- Word limit: ≤ 2000 words
- ≤ 15 references
- ≤ 3 figures/graphs/pictures

Perspectives

An unstructured article discussing an issue regarding the surveillance of and response to public health events. The scope of the discussion must be clearly defined.

- Word limit: ≤ 1000 words
- ≤ 10 references
- ≤ 1 illustration

Case Report or Case Series

An unstructured article describing an unusual case or series of cases of public health significance. Subheadings may be used to increase the readability of the article.

- Unstructured abstract of ≤ 250 words
- Word limit: ≤ 2000 words
- ≤ 15 references
- ≤ 3 figures/graphs/pictures

Regional Analysis

An article providing an analysis of a topic for the Western Pacific Region, typically authored by WHO staff as part of their routine work on behalf of Member States. Regional Analyses do not undergo peer review.

Letter to the Editor

A letter commenting on a previously published article OR a letter commenting on the theme of the issue. Letters do not undergo peer review.

- Word limit: ≤ 500 words
- ≤ 5 references
- ≤ 1 illustration

News, Meeting and Conference Reports

News items and meeting and conference reports do not undergo peer review. Please contact the Coordinating Editor at WPSAR@wpro.who.int if you intend on submitting such an article.

Illustrations

Refer to the article type for the limit on illustrations (figures/graphs/pictures). Please insert all illustrations at the end of the article with titles. Each illustration must be referred to in the text and must be understood on its own. Use Microsoft® Office Excel for graphs and Microsoft® Office Word for tables and diagrams. Additionally, please provide a Microsoft® Office Excel spreadsheet of the data used to create a graph. Footnotes should be placed under the illustration and should use the following symbols in superscript format: *, †, ‡, §, ||, **, ††, etc.

References

Reference the most recent and relevant publications. Please use the Vancouver referencing style with in-text citations and a bibliography at the end of the text. Sample references can be viewed on the National Institutes of Health website.

Place the bibliography at the end of the article text and not as footnotes. Write journal names in full. Use superscript sequential numbering for citing references in the text. Place the number after any punctuation. For example:

These results are consistent with the original study.¹¹

Reference personal communication in the text only and include the person's full name and institution.

Caution should be used in referencing websites; it should be done only when their content has been substantially described in the article.

Peer review process

Every article is initially screened by the Editorial Team to ensure it fits the scope of the journal. All articles, with the exception of regional analyses, letters to the editor, news items and meeting and conference reports, then undergo external peer review by two reviewers. This blind peer review process ensures that the reviewer does not know the identity of the author(s) and the author(s) do not know the identity of the reviewer. Significant effort is made to make this process timely, but since it relies on the availability and cooperation of persons external to the journal, it can take considerable time.

Upon receipt of the reviews, the Coordinating Editor assesses the comments and recommendations made by the reviewers, and then decides on the outcome of the peer review process. One of four options will be chosen: accept submission, accept with revisions, submit for review, or decline submission. The corresponding author will be advised of this outcome.

If the article has been accepted or accepted with revisions are required, you will be invited to revise your article according to the reviewer comments. A separate MS Word document outlining how you addressed each of the reviewer comments is also required. You must indicate the page and paragraph numbers where the changes were made and should provide reasons for not making a suggested change. Both the changes and reasons will be assessed

against the reviewer comments by the Coordinating Editor and may require further clarification from the authors. Once all comments have been adequately addressed, the article will commence the publication process.

If the outcome of the review process is “submit for review”, then the same process is followed. However, the resubmitted article and responses to the reviewer comments are sent back to the original reviewers for another round of peer review. You will be asked to respond to a second round of reviewer comments, which will again be assessed by the Coordinating Editor. Once both sets of reviewer comments have been adequately addressed, the article will commence the publication process.

The publication process comprises rigorous editing for content and style by an external technical editor, followed by layout and proofreading. Authors may be asked to provide further information or clarifications during these stages. An article is not formally accepted for publication until these stages have been completed and approval has been granted by the Editorial Team. The authors will also have an opportunity to approve the final proof prior to publication on the WPSAR website. The article will be batched with others in the next quarterly issue.

Authorship

As per the International Committee of Medical Journal Editors (ICMJE), all authors should have contributed significantly to the article through one or more of the following in each category A, B and C:

A

- Study design
- Data collection
- Data analysis
- Data interpretation

B

- Drafting the article
- Critically revising the article

C

- Final approval of the article for submission

Any other contributors may be listed in the Acknowledgements section.

Acknowledgements

Contributors who do not fulfil the authorship requirements may be acknowledged. Permission from all contributors in the acknowledgement section should be sought. We assume that permission has been granted and will not follow up with the authors to confirm.

Ethics and permissions

It is the responsibility of authors to gain appropriate ethics approval for their work. A statement of ethics approval obtained or an explanation of why ethics approval was not required should be included for all articles during the submission process.

License for publication

Prior to publication, all authors are asked to sign a licence to publish that grants exclusive copyright license to the World Health Organization.

Conflicts of interest

A conflict of interest is defined by ICMJE as “when an author or author’s institution, reviewer, or editor has financial or personal relationships that inappropriately influence (bias) his or her actions”. Conflicts of interest may be financial, institutional, research or personal. A relationship does not always represent a conflict of interest and does not necessarily preclude publication in WPSAR. All authors and reviewers will be required to state any potential conflicts of interest, which will be assessed by the Editorial Team.

Funding

Authors will be required to state the sources of funding for their work.

Photographs for cover

If authors have taken photographs that are relevant to their article, they may be submitted for consideration for publication on the cover of the issue. Submission of a photograph does not guarantee its publication.

Language

Articles should be written in English. Authors who require assistance with preparing their articles in English should contact WPSAR at WPSAR@wpro.who.int. Once published, all abstracts and most articles are translated into Chinese.

Article submission process

Submit articles to the Coordinating Editor through the online journal management system on the WPSAR website. When submitting the article, you will be requested to provide the following:

- a cover letter describing the article and why it should be published;
- a title page with:
 - the article title,
 - a short title,
 - a brief description of the article of ≤ 50 words,
 - ≤ 7 keywords,
 - full names of all authors and institutions,
 - full contact details of the corresponding author,
 - data in an MS Excel spreadsheet for any graphs
 - names and e-mail addresses of two suggested reviewers (optional but recommended);
- acknowledgements, conflicts of interest, ethics statement and funding information (attached as a separate file to ensure a blind review);
- an MS Word file or equivalent of the article; and
- a scanned copy of the WPSAR licence for publication signed by all authors.

With the online journal management system, you will be able to track the progress of your article through the editorial process. If you encounter any difficulties with this system, please refer to our *WPSAR online journal system – User guide for authors*.

Corrections

If authors of a published article become aware of any errors with the article, they should contact the Coordinating Editor at WPSAR@wpro.who.int. Corrections will be published online.

Acknowledgement to WPSAR reviewers 2013

The WPSAR Editorial Team is grateful to all reviewers that have contributed their time and expertise to peer-review our articles. We thank them for their support. Our apologies to those reviewers we may have inadvertently missed.

Our reviewers for articles published in 2013 were:

Bijay Adhikari, Martha Anker, Christian Auer, Aridam Basu, Cynthia Chee, Siddhartha Sankar Datta, Stephanie Davis, Jose Derraik, Maria Nerissa Dominguez, Georgina Dove, Marion Easton, Keith Eastwood, Marsha L Feske, Emma Jane Field, James Fielding, Simon Firestone, Alice Ruth Foxwell, Philippe Glaziou, Richard James Hall, Max Hardiman, Mike Kama, Kamal Kishore, Chia-Hsien Lin, Constance Low, SH Lum, John S Mackenzie, Peter D Massey, Lisa McCallum, Ellen Mitchell, Rodney Moran, Keiko Nakamura, Lee-Ching Ng, Ni Daxin, Akihiro Ohkado, Kosuke Okada, Amy Elizabeth Parry, Beverly Paterson, Z Qiaoli, John Rainford, Maria Conception Rey Rocas, John Stanley Rule, Mika Saito, Marcel Salanthe, Gina Samaan, Saraswathi Bina Rai, Jacques Sebert, Shi Lizheng, Shoji Yoshimatsu, Shuko Nagai, Kevin Soli, Pawel Stefanoff, Motoi Suzuki, A Tamaru, Betsy Todd, Daniel Michael Tompkins, Huu Dat Tran, Raman Velayudhan, Polly Wallace, Wang Xu, Xing Jun, Dongbao Yu, Aysha Zahidie, Weigong Zhang



wpsar@wpro.who.int | www.wpro.who.int/wpsar